37 mm

37 mm

Total bilirubi

utrophils

the normal range [see Warnings and Precautions (5.13)].

convulsions, hypoesthesia, paresthesia, neuropathy, tremor, vertigo.

Liver and Biliary System: hepatic enzyme increase, hepatic failure, hepatitis

 $\textit{Skin and Appendages}: erythema\ multiforme, photoallergic\ dermatitis, Stevens-Johnson\ syndrome and photoallergic\ dermatitis, and photoallergic\ dermatitis\ dermatitis\$ 

Body as a Whole: redistribution/accumulation of body fat [see Warnings and Precautions (5.15)].

Hemic and Lymphatic: anemia (including pure red cell aplasia and severe anemias progressing on therapy).

Hepatobiliary Disorders: hepatic steatosis, hepatitis, increased liver enzymes (most commonly AST, ALT gamma GT).

rhabdomyolysis, osteomalacia, hypokalemia, muscular weakness, myopathy, hypophosphatemia.

nistered with other antiretroviral medications for treatment of HIV-1 infection

[see Warnings and Precautions (5.4)]. Drugs that decrease renal function may increase concent

EFV\*

Lanticonvulsant LEFV

Lbupropion'

sertraline

Lhydroxyitraconazole<sup>3</sup>

posaconazole

clarithromycin\*

rifabutin\*

EFV\*

14-OH metabolite\*

dihydroartemisinin

desacetyl diltiazem\*

calcium channel block

atovaguone

atorvastatin'

simvastatin

boceprevir\*

elbasvir

pibrentasvi

, simeprevir\* → EFV

velpatasv

active metabolites of

immunosuppressant

\* The interaction between EFV and the drug was evaluated in a clinical study. All other drug interactions shown are predicted

etonogestrel

methadone

TDF

proguanil

 ${\it Metabolic and Nutritional:} \ hypercholester olemia, hypertrigly ceridemia.$ 

Musculoskeletal: muscle weakness, CPK elevation, rhabdomyolvsis,

 ${\it Immune System \, Disorders:} \ all ergic \, reaction, including \, angioedema.$ 

Gastrointestinal Disorders: pancreatitis, increased amylase, abdominal pain.

Respiratory, Thoracic, and Mediastinal Disorders: dyspnea.

General Disorders and Administration Site Conditions: asthenia.

 $7.1 \quad Not\,Recommended\,with\,Other\,Antiretroviral\,Medications$ 

polyuria [see Warnings and Precautions (5.4)].

Skin and Subcutaneous Tissue Disorders: rash.

fractures), muscular weakness, myopathy.

drug with a known risk of Torsade de Pointes

7.3 Drugs Affecting Renal Function

7.4 Cannabinoid Test Interaction

Isee Clinical Pharmacology (12.3)1.

plasma concentrations.

Warfarin

Phenytoin

Carbamazepine

Phenobarbital

itidepressants

Bupropion

Sertraline

Intifungals:

Ketoconazole

Posaconazole

Clarithromycir

Antimycobacteria

Rifampin

Atovaguone

proguanil

Diltiazem

Artemether/lumefantrine

Calcium channel blockers

Others (e.g., felodipine

nicardipine, nifedipine,

lepatitis C antiviral agents

Elbasvir/Grazoprevir

Pibrentasvir/Glecaprevir

Velpatasvir/Sofosbuvir/Voxilanre

Ledipasvir/Sofosbuvi

Hepatitis B antiviral agents

Adefovir dipivoxil

Hormonal contraceptives

Oral Ethinyl estradiol

Norgestimate Implant Etonogestrel

Cyclosporine, tacrolimus,

sirolimus, and others

metabolized by CYP3A

Narcotic analgesic

This table is not all-inclusive

7.6 Drugs without Clinically Significant Interactions

Methadone

HMG-CoA reductas

Atorvastatir

Simvastatin

Boceprevir

nhibitors

Anti-infective:

concentrations when coadministered with EFV.

les potentially significant interactions, but is not all incl

Based on Drug Interaction Studies or Predicted Interactio

Concomitant Drug Class: Drug Name | Effect

DRUG INTERACTIONS

7.2 QT Prolonging Drugs

 ${\it Metabolism and Nutrition Disorders:} lactic acidosis, hypokalemia, hypophosphatemia. \\$ 

cointestinal: constipation, malabsorption

Musculoskeletal: arthralgia, myalgia, myopathy.

Special Senses: abnormal vision, tinnitus.

Endocrine and Metabolic: hyperglycemia.

Hypersensitivity: anaphylaxis, urticaria.

Cardiovascular: flushing, palpitations.

6.2 Postmarketing Experience

Endocrine: gynecomastia.

Respiratory: dyspnea.

Lamivudine

General: weakness.

Precautions (5.1, 5.2)1.

Skin: Alopecia, pruritus.

Tenofovir Disoproxil Fumarate

Fifty-three percent (531/1 008) of nationts receiving FFV a component of efavirenz lamivudine and tenofovir disproxil furnarate tablets in

controlled trials reported central nervous system symptoms (any grade, regardless of causality) compared to 25% (156/635) of patients receiving control regimens. These symptoms included, but were not limited to, dizziness (28.1% of the 1,008 patients), insomnia (16.3%), impaired

and 2.1% of patients discontinued therapy as a result. These symptoms usually begin during the first or second day of therapy and generally resolve

after the first 2 to 4 weeks of therapy. After 4 weeks of therapy, the prevalence of nervous system symptoms of at least moderate severity ranged from 5% to 9% in patients treated with regimens containing EFV and from 3% to 5% in patients treated with a control regimen. Inform patients that

these common symptoms were likely to improve with continued therapy and were not predictive of subsequent onset of the less frequent psychiatric symptoms [see Warnings and Precautions (5.5)]. Dosing at bedtime may improve the tolerability of these nervous system symptoms

In the ENCORE1 study, at Week 48, 40% of EFV 400 mg recipients and 48% of EFV 600 mg recipients reported central nervous system disorders. The most common symptoms (> 10%) were dizziness (27% vs. 35%) and headache (11% vs. 11%).

Late-onset neurotoxicity, including ataxia and encephalopathy (impaired consciousness, confusion, psychomotor slowing, psychosis, delirium),

may occur months to years after beginning efavirenz therapy. Some events of late-onset neurotoxicity have occurred in patients with CYP2B6 genetic polymorphisms which are associated with increased efavirenz levels despite daily dosages of 600 mg of efavirenz. Patients presenting with

igns and symptoms of serious neurologic adverse experiences should be evaluated promptly to assess the possibility that these events may be

EFV, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, may cause fetal harm when administered during the firs

trimester to a pregnant woman. Advise females of reproductive potential who are receiving EFV to avoid pregnancy [see Use in Specific Populations

of patients treated in control groups. Rash associated with blistering, moist desguamation, or ulceration occurred in 0.9% (9/1,008) of patients

treated with EFV. The incidence of Grade 4 rash (e.g., erythema multiforme, Stevens-Johnson syndrome) in patients treated with EFV in all studies and expanded access was 0.1%. Rashes are usually mild-to-moderate maculopapular skin eruptions that occur within the first 2 weeks of initiating

therapy with EFV (median time to onset of rash in adults was 11 days) and, in most patients continuing therapy with EFV, rash resolves within 1

EFV can generally be reinitiated in patients interrupting therapy because of rash. EFV should be discontinued in patients developing severe rash

associated with blistering, desquamation, mucosal involvement, or fever. Appropriate antihistamines and/or corticosteroids may improve the tolerability and hasten the resolution of rash. For patients who have had a life-threatening cutaneous reaction (e.g., Stevens-Johnson syndrome),

In the ENCORE1 study at Week 48, different types of rash (such as rash, rash papular, rash maculopapular and rash pruritic) occurred in 32% of EFV

recipients. The discontinuation rate for rash in the ENCORE1 study was 3% of EFV 600 mg recipients and 1% of EFV 400 mg recipients

600 mg recipients and 26% of EFV 400 mg recipients. Grade 3 to 4 rash was reported in 3% of EFV 600 mg recipients and 1% of EFV 400 mg

tmarketing cases of hepatitis, including fulminant hepatitis progressing to liver failure requiring transplantation or resulting in death, have beer

reported in patients treated with EFV. Reports have included patients with underlying hepatic disease, including coinfection with hepatitis B or C,

EFV, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, is not recommended for patients with moderate or severe

Monitoring of liver enzymes before and during treatment is recommended for all patients [see Dosage and Administration (2.1)]. Consider

Discontinue efavirenz, lamivudine and tenofovir disoproxil fumarate if elevation of serum transaminases is accompanied by clinical signs or

In pediatric patients with a history of prior antiretroyiral nucleoside exposure, a history of pancreatitis, or other significant risk factors for the

development of pancreatitis, 3TC, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, should be used with caution.

Treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate should be stopped immediately if clinical signs, symptoms, or laboratory

Toxicology (13.2)]. Caution should be taken in any patient with a history of seizures. Patients who are receiving concomitant anticonvulsant

medications primarily metabolized by the liver, such as phenytoin and phenobarbital, may require periodic monitoring of plasma levels [see Drug

Treatment with EFV has resulted in increases in the concentration of total cholesterol and triglycerides. Cholesterol and triglyceride testing should

biochemical markers of bone metabolism, suggesting increased bone turnover relative to comparators [see Adverse Reactions (6.1)]. Serum

The effects of TDF-associated changes in BMD and biochemical markers on long-term bone health and future fracture risk in adults and pediatric

subjects 2 years and older are unknown. The long-term effect of lower spine and total body BMD on skeletal growth in pediatric patients, and in

Although the effect of supplementation with calcium and vitamin D was not studied, such supplementation may be beneficial for all. Assessment of

BMD should be considered for adult and pediatric patients who have a history of pathologic bone fracture or other risk factors for osteoporosis or

Mineralization Defects: Cases of osteomalacia associated with proximal renal tubulopathy, manifested as bone pain or pain in extremities and

which may contribute to fractures, have been reported in association with TDF use (see Adverse Reactions (6.2)). Arthralgia and muscle pain or

weakness have also been reported in cases of proximal renal tubulopathy. Hypophosphatemia and osteomalacia secondary to proximal renal

tubulopathy should be considered in patients at risk of renal dysfunction who present with persistent or worsening bone or muscle symptoms while

and TDF. During the initial phase of combination antiretroviral treatment, patients whose immune system responds may develop an inflammator

response to indolent or residual opportunistic infections (such as Mycobacterium avium infection, cytomegalovirus, Pneumocystis jirovecii

Autoimmune disorders (such as Graves' disease, polymyositis, Guillain-Barre syndrome, and autoimmune hepatitis) have also been reported to

occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

In HIV-infected nations, redistribution/accumulation of body fat including central obesity dorsocervical fat enlargement (buffalo hump)

antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been

QTc prolongation has been observed with the use of EFV [see Drug Interactions (7.2, 7.5) and Clinical Pharmacology (12.2)]. Consider alternatives

to products containing EFV when coadministered with a drug with a known risk of Torsade de Pointes or when administered to patients at higher

Because clinical studies are conducted under widely varying conditions, the adverse reaction rates observed in clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

In Trial 903, 600 antiretroviral-naïve subjects received TDF (N = 299) or stavudine (d4T) (N = 301) administered in combination with 3TC and EFV

for 144 weeks. The most common adverse reactions were mild to moderate dastrointestinal events and dizziness. Mild adverse reactions (Grade 1)

were common with a similar incidence in both arms and included dizziness, diarrhea, and nausea. Table 1 provides the treatment-emergent adverse

d4T + 3TC + EFV

N = 301

12%

17%

12%

13%

TDF + 3TC + EFV

14%

13%

11%

N = 299

econstitution syndrome has been reported in HIV-infected patients treated with combination antiretroviral therapy, including EFV, 3TC,

Bone Mineral Density (BMD): In clinical trials in HIV-1-infected adults, TDF was associated with slightly greater decreases in BMD and increases in BMD and increase in BMD and increases in BMD and increase in BMD and in

ions have been observed in patients receiving EFV, generally in the presence of known medical history of seizures [see Nonclinical

ic impairment. Careful monitoring is recommended for patients with mild hepatic impairment receiving EFV [see Adverse Reactions (6.1) and

ntinuing efavirenz, lamivudine and tenofovir disoproxil fumarate in patients with persistent elevations of serum transaminases to greater than

trolled clinical trials, 26% (266/1,008) of patients treated with 600 mg EFV experienced new-onset skin rash compared with 17% (111/635)

elated to efavirenz use, and whether discontinuation of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets is warranted

month (median duration, 16 days). The discontinuation rate for rash in clinical trials was 1.7% (17/1.008).

[see Dosage and Administration (2.2)].

5.7 Embryo-Fetal Toxicity

Use in Specific Populations (8.7)].

5.11 Convulsions

Interactions (7.5)].

Revised: 08/2023

5.12 Lipid Elevations

five times the upper limit of the normal range.

5.13 Bone Loss and Mineralization Effects

5.14 Immune Reconstitution Syndrome

5.16 QTc Prolongation

risk of Torsade de Pointe

ADVERSE REACTIONS

6.1 Clinical Trials Experience

Rash event

Headache

Diarrhea

Back pain

mptoms of hepatitis or hepatic decompensation

5.8 Skin and Systemic Hypersensitivity Reaction

alternate therapy should be considered [see Contraindications (4)].

and patients without pre-existing hepatic disease or other identifiable risk factors.

abnormalities suggestive of pancreatitis occur [see Adverse Reactions (6.1)].

be performed before initiating EFV therapy and at periodic intervals during therapy.

particular, the effects of long-duration exposure in younger children is unknown.

receiving TDF-containing products [see Warnings and Precautions (5.4)].

peripheral wasting, facial wasting, breast enlargement, and

The following adverse reactions are discussed in other sections of the labeling:

Psychiatric Symptoms [see Warnings and Precautions (5.5)].

Hepatotoxicity [see Warnings and Precautions (5.9)].

Fat Redistribution [see Warnings and Precautions (5.15)].

Pancreatitis [see Warnings and Precautions (5.10)].

Ffavirenz | Lamivudine and Tenofovir Disonroxil Fumarate

Clinical Trials in Treatment-Naïve HIV-1 Infected Adult Subjects

Nervous System Symptoms [see Warnings and Precautions (5.6)].

Exacerbations of Hepatitis B [see Boxed Warning, Warnings and Precautions (5.1)].

New Onset or Worsening Renal Impairment [see Warnings and Precautions (5.4)].

Skin and Systemic Hypersensitivity Reaction [see Warnings and Precautions (5.8)].

Bone Loss and Mineralization Effects [see Warnings and Precautions (5.13)].

Immune Reconstitution Syndrome [see Warnings and Precautions (5.14)].

Lactic Acidosis/Severe Hepatomegaly with Steatosis [see Warnings and Precautions (5.2)].

reactions (Grades 2 to 4) occurring in greater than or equal to 5% of subjects treated in any treatment group

Table 1. Selected Adverse Reactions³ (Grades 2 to 4) Reported in ≥5% in Any Treatment Group in Trial 903 (0 to 144 Weeks)

parathyroid hormone levels and 1,25 Vitamin D levels were also higher in subjects receiving TDF.

 $bone\ loss.\ If\ bone\ abnormalities\ are\ suspected\ then\ appropriate\ consultation\ should\ be\ obtained.$ 

pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment.

tion (8.3%), somnolence (7.0%), abnormal dreams (6.2%), and hallucinations (1.2%). These symptoms were severe in 2.0% of patients

receiving 3TC alone or in combination with other antiretroviral agents [see Warnings and Precautions (5.10)].

Pancreatitis: Pancreatitis, which has been fatal in some cases, has been observed in antiretroviral nucleoside-experienced pediatric subjects

Changes in Bone Mineral Density: In HIV-1-infected adult subjects in Trial 903, there was a significantly greater mean percentage decrease from baseline in BMD at the lumbar spine in subjects receiving TDF + 3TC + EFV (-2.2% ± 3.9) compared with subjects receiving d4T + 3TC + EFV (-1.0%

 $\pm$  4.6) through 144 weeks. Changes in BMD at the hip were similar between the two treatment groups (-2.8%  $\pm$  3.5 in the TDF group vs. -2.4%  $\pm$  4.5 in the d4T group). In both groups, the majority of the reduction in BMD occurred in the first 24 to 48 weeks of the trial and this reduction was

sustained through Week 144. Twenty-eight percent of TDF-treated subjects vs. 21% of the d4T-treated subjects lost at least 5% of BMD at the spine or 7% of BMD at the hip. Clinically relevant fractures (excluding fingers and toes) were reported in 4 subjects in the TDF group and 6 subjects in the

d4T group. In addition, there were significant increases in biochemical markers of bone metabolism (serum bone-specific alkaline phosphatase, serum osteocalcin, serum C telopeptide, and urinary N telopeptide) and higher serum parathyroid hormone levels and 1,25 Vitamin D levels in the

 $TDF group\ relative\ to\ the\ d4T\ group; however, except\ for\ bone-specific\ alkaline\ phosphatase,\ these\ changes\ resulted\ in\ values\ that\ remained\ within\ the phosphatase,\ the properties of the phosphatase and the phosphatase are changed by the properties of the phosphatase and the phosphatase are changed by the phosphatase ar$ 

e following adverse reactions have been identified during post-approval use for each of the individual components of efavirenz, lamivudine and

tenofovir disoproxil fumarate tablets (EFV. 3TC, and TDF). Because these reactions are reported voluntarily from a population of unknown size, it is

not always possible to reliably estimate their frequency or establish causal relationship to drug exposure. These reactions have been chosen for

Central and Peripheral Nervous System: abnormal coordination, ataxia, encephalopathy, cerebellar coordination and balance disturbances,

Hepatic and Pancreatic: lactic acidosis and hepatic steatosis, posttreatment exacerbation of hepatitis B [see Boxed Warning, Warnings and

Renal and Urinary Disorders: renal insufficiency, acute renal failure, renal failure, acute tubular necrosis, Fanconi syndrome, proximal renal

tubulopathy, interstitial nephritis (including acute cases), nephrogenic diabetes insipidus, renal insufficiency, increased creatinine, proteinuria,

Musculoskeletal and Connective Tissue Disorders: rhabdomyolysis, osteomalacia (manifested as bone pain and which may contribute to

The following adverse reactions, listed under the body system headings above, may occur as a consequence of proximal renal tubulopathy.

Efavirenz, lamivudine and tenofovir disoproxil fumarate is a complete regimen for the treatment of HIV-1 infection; therefore, it should not be

There is limited information available on the potential for a pharmacodynamic interaction between EFV and drugs that prolong the QTc interval. QTc

Tenofovir is primarily eliminated by the kidneys [see Clinical Pharmacology (12.3)]. Coadministration of EFV/3TC/TDF with drugs that are

eliminated by active tubular secretion may increase concentrations of tenofovir and/or the coadministered drugs. Some examples include, but are not limited to, acyclovir, cidofovir, ganciclovir, valganciclovir, aminoglycosides (e.g., gentamicin), and high-dose or multiple NSAIDs

EFV does not bind to cannabinoid receptors. False-positive urine cannabinoid test results have been reported with some screening assays in

uninfected and HIV-infected subjects receiving EFV. Confirmation of positive screening tests for cannabinoids by a more specific method is

EFV has been shown in vivo to induce CYP3A and CYP2B6. Other compounds that are substrates of CYP3A or CYP2B6 may have decreased plasma

that induce CYP3A activity (e.g., phenobarbital, rifampin, rifabutin) would be expected to increase the clearance of EFV resulting

No drug interaction studies have been conducted using efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. However, drug interaction

Clinical Comment

nitor INR and adjust warfarin dosage if necessary.

Alternative anticonvulsant treatment should be used.

for itraconazole or ketoconazole can be made.

ifampin to patients weighing 50 kg or more

Concomitant administration is not

the full prescribing information for diltiazer

oncomitant administration of boceprevir

sponse to elbasvir/grazoprevir.

sofosbuvir/velpatasvir.

effect of sofosbuvir/velpatasvir

ormonal contraceptives

treatment with EFV.

with etonogestrel in EFV-exposed patients.

recommended.

interval prolongation

ere are insufficient data to make a dose recommendation for EFV

Monitor anticonvulsant plasma levels periodically because of potential for reduction in anticonvulsant and/or EFV plasma levels.

Increases in bupropion dosage should be guided by clinical response

upropion dose should not exceed the maximum recommended dos

creases in sertraline dosage should be guided by clinical response.

void concomitant use unless the benefit outweighs the risks.

crease EFV total daily dose to 800 mg once daily when coadr

nterval prolongation [see Warnings and Precautions (5.16)].

onsider alternative antifungal treatment because no dose recommendation

onsider alternatives to macrolide antibiotics because of the risk of QT

crease daily dose of rifabutin by 50%. Consider doubling the rifabutin dos

Diltiazem dose adjustments should be guided by clinical response (refer to

When coadministered with EFV, dosage adjustment of calcium channel

(refer to the full prescribing information for the calcium channel blocker).

blocker may be needed and should be guided by clinical response

Plasma concentrations of atorvastatin, pravastatin, and simvastatir decreased. Consult the full prescribing information for the HMG-CoA reductase inhibitor for guidance on individualizing the dose.

padministration of EFV with elbasvir/grazoprevir is contraindicated

padministration of EFV is not recommended because it may lead

oncomitant administration of simeprevir is not recommended.

recommended because it may result in loss of therapeutic effect of

padministration of EFV and sofosbuvir/velpatasvir/voxilaprevir is

oncomitant administration of adefovir dipivoxil is not recommended.

A reliable method of barrier contraception should be used in addition to

A reliable method of barrier contraception should be used in addition to

ormonal contraceptives. Decreased exposure of etonogestrel may be expected. There have been postmarketing reports of contraceptive failure

Dose adjustments of the immunosuppressant may be required. Close

monitoring of immunosuppressant concentrations for at least 2 weeks (until stable concentrations are reached) is recommended when starting or stopping

Monitor for signs of methadone withdrawal and increase methadone

not recommended because it may result in loss of therapeutic

Monitor for adverse reactions associated with TDF.

Coadministration of EFV and sofosbuvir/velpatasvir is not

to reduced therapeutic effect of pibrentasvir/glecaprevir.

[see Contraindications (4)] because it may lead to loss of virologic

prolongation has been observed with the use of EFV [see Clinical Pharmacology (12.2)]. Consider alternatives to EFV when coa

inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to EFV, 3TC, and TDF.

Body as a Whole: alleroic reactions, asthenia, redistribution/accumulation of body fat [see Warnings and Precautions (5.15)].

Psychiatric: aggressive reactions, agitation, delusions, emotional lability, mania, neurosis, paranoia, psychosis, suicide, catatoni

evaluated in a limited number of women as reported to the APR. Available data from the APR show no difference in the risk of major birth defects for EFV and 3TC compared to the background rate for major birth defects of 2.7% in the U.S. reference population of the Metropolitan Atlanta Congenital Defects Program (MACDP). Available data from the APR also show no increase in the overall risk of major birth defects with first trimester exposure for TDF (2.1%) compared with the background rate for major birth defects of 2.7% in a U.S. reference population of the MACDF (see Data). 3TC produced embryonic toxicity in rabbits at a dose that produced similar human exposures as the recommended clinical dose. The relevance of animal findings to human pregnancy registry data is not known. The rate of miscarriage is not reported in the APR. The estimated background rate of miscarriage in clinically recognized pregnancies in the U.S. general population is 15% to 20%. The background risk for major birth defects and miscarriage for the indicated population is unknown. The APR

uses the MACDP as the U.S. reference population for birth defects in the general population. The MACDP evaluates women and infants from a limited geographic area and does not include outcomes for births that occurred at less than 20 weeks' gestation Human Data: Efavirenz: There are retrospective postmarketing reports of findings consistent with neural tube defects, including meningomyelocele, all in infants of mothers exposed to EFV- containing regimens in the first trimester [see Warnings and Precautions (5.7)]. Based on prospective reports from the APR of approximately 1,000 live births following exposure to EFV-containing regimens (including over 800 live births exposed in the first trimester), there was no difference between EFV and overall birth defects compared with the background birth defect rate of 2.7% in the U.S. reference population of the Metropolitan Atlanta Congenital Defects Program. As of the interim APR report issued December

anophthalmia. Lamivudine: Based on prospective reports from the APR of over 11,000 exposures to 3TC during pregnancy resulting in live births (including over 4,300 exposed in the first trimester), there was no difference between 3TC and overall risk of birth defects for 3TC compared with the background birth defect rate of 2.7% in the U.S. reference population of the MACDP. The prevalence of defects in live births was 3.1% (95% CI: 2.6% to 3.6%) following first trimester exposure to 3TC-containing regimens and 2.8% (95% CI: 2.5% to 3.3%) following second/third trimester exposure to 3TC containing regimens. 3TC pharmacokinetics were studied in pregnant women during 2 clinical trials conducted in South Africa. The trials assessed pharmacokinetics in 16 women at 36 weeks' gestation using 150 mg 3TC twice daily with zidovudine, 10 women at 38 weeks' gestation using 150 mg 3TC twice daily with zidovudine, and 10 women at 38 weeks' gestation using 3TC 300 mg twice daily without other antiretrovirals. These trials were not designed or

3TC concentrations were generally similar in maternal peopatal and umbilical cord serum samples. In a subset of subjects, amniotic fluid

specimens were collected following natural rupture of membranes and confirmed that 3TC crosses the placenta in humans. Based on limited data at

delivery, median (range) amniotic fluid concentrations of 3TC were 3.9 (1.2 to 12.8)-fold greater compared with paired maternal serum

powered to provide efficacy information.

2014, the prevalence of birth defects following first-trimester exposure was 2.3% (95% CI: 1.4% to 3.6%). One of these prospectively reported defects with first-trimester exposure was a neural tube defect. A single case of anophthalmia with first-trimester exposure to EFV has also been

prospectively reported. This case also included severe oblique facial clefts and amniotic banding, which have a known association with

live births (including 3.342 exposed in the first trimester and 1.475 exposed in the second/third trimester), there was no increase in overall major birth defects with TDF compared with the background birth defect rate of 2.7% in a U.S. reference population of the MACDP. The prevaled maior birth defects in live births was 2.3% (95% CI: 1.8% to 2.8%) with first trimester exposure to TDF-containing regimens, and 2.1% (95% CI: 1.4% to 3.0%) with the second/third trimester exposure to TDF-containing regimens.Prospective reports from the APR of overall major birth defects in pregnancies exposed to TDF are compared with a U.S. background major birth defect rate. Methodological limitations of the APR include the use of MACDP as the external comparator group. Limitations of using an external comparator include differences in methodology and populations, as well as confounding due to the underlying disease.

In published data from three controlled clinical trials, a total of 327 pregnant women with chronic HBV infection were administered tenofovir disoproxil fumarate from 28 to 32 weeks gestation through 1 to 2 months postpartum and followed for up to 12 months after delivery. There were no new safety findings in pregnant women compared with the known safety profile of tenofovir disoproxil fumarate in HBV-infected adults. An increased risk of adverse pregnancy-related outcomes was not observed; 2 stillbirths were identified, and there was 1 major birth defect (talipes) and 1 occurrence of multiple congenital abnormalities (not further specified) in tenofovir disoproxil furnarate-exposed infants. Infants were followed for up to 12 months after delivery; there were no clinically relevant drug-related safety findings in infants exposed to tenofovir disoproxil fumarate during late gestation Animal Data: Efavirenz: Effects of EFV on embryo-fetal development have been studied in three nonclinical species (cynomolgus monkeys, rats,

and rabbits). In monkeys, EFV 60 mg/kg/day was administered to pregnant females throughout pregnancy (gestation days 20 through 150). The maternal systemic drug exposures (AUC) were 1.3 times the exposure in humans at the recommended clinical dose (600 mg/day), with fetal umbilical venous drug concentrations approximately 0.7 times the maternal values. Three of 20 fetuses/infants had one or more malformations: there were no malformed fetuses or infants from placebo-treated mothers. The malformations that occurred in these three monkey fetuses included anencephaly and unilateral anophthalmia in one fetus, microophthalmia in a second, and cleft palate in the third. There was no NOAEL (no observable adverse effect level) established for this study because only one dosage was evaluated. In rats, EFV was administered either during organogenesis (gestation days 7 to 18) or from gestation day 7 through lactation day 21 at 50, 100, or 200 mg/kg/day. Administration of 200 mg/kg/day in rats was associated with increase in the incidence of early resorptions; and doses 100 mg/kg/day and greater were associated with early neonatal mortality. The AUC at the NOAEL (50 mg/kg/day) in this rat study was 0.1 times that in humans at the recommended clinical dose. Drug concentrations in the milk on lactation day 10 were approximately 8 times higher than those in maternal plasma. In pregnant rabbits, EFV was neither embryo lethal nor teratogenic when administered at doses of 25, 50, and 75 mg/kg/day over the period of organogenesis (gestation days 6  $through 18). The AUC at the NOAEL (75\,mg/kg/day) in rabbits was 0.4 times that in humans at the recommended clinical dose and the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the commended clinical dose and the commended clinical dose are the clinical dose are the clinical dose are the clinical dose$ Lamivudine: 3TC was administered orally to pregnant rats (at 90, 600, and 4,000 mg per kg per day) and rabbits (at 90, 300, and 1,000 mg per kg per day and at 15, 40, and 90 mg per kg per day) during organogenesis (on gestation Days 7 through 16 [rat] and 8 through 20 [rabbit]). No evidence of fetal malformations due to 3TC was observed in rats and rabbits at doses producing plasma concentrations (C<sub>mm</sub>) approximately 35 times higher than human exposure at the recommended daily dose. Evidence of early embryolethality was seen in the rabbit at system exposures (AUC) similar to those observed in humans, but there was no indication of this effect in the rat at plasma concentrations  $(C_{max})$  35 times higher than human exposure at the recommended daily dose. Studies in pregnant rats showed that 3TC is transferred to the fetus through the placenta. In the fertility/pre- and postnatal development study in rats, 3TC was administered orally at doses of 180, 900, and 4,000 mg per kg per day (from prior to mating through postnatal Day 20). In the study, development of the offspring, including fertility and reproductive performance, was not affected by maternal administration of 3TC Tenofovir Disoproxil Furnarate: TDF was administered orally to pregnant rats (at 0, 50, 150, or 450 mg/kg/day) and rabbits (at 0, 30, 100, or 300 mg/kg/day) through organogenesis (on gestation days 7 through 17, and 6 through 18, respectively). No significant toxicological effects were

observed in embryo-fetal toxicity studies performed with TDF in rats at doses up to 14 times the human dose based on body surface area comparisons and in rabbits at doses up to 19 times the human dose based on body surface area comparisons. In a pre/postnatal development study  $in \ rats, TDF \ was administered \ or ally through lactation \ at doses \ up \ to 600 \ mg/kg/day; \ no \ adverse \ effects \ were \ observed \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ tenofovir \ in \ the \ offspring \ at \ at \ the \ offspring \ at \ offspring \ at \ the \ offspring \ at \ offspring \ at \ \ offspring \ at \ offspring \ a$ exposures of approximately 2.7 times higher than human exposures at the recommended daily dose of TD Risk Summary: The Centers for Disease Control and Prevention recommend that HIV-1-infected mothers not breastfeed their infants to avoid

risking postnatal transmission of HIV-1 infection. effects of EFV on milk production. s have been conducted with the individual components of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets (EFV, 3TC, and TDF)

on the effects of 3TC on the breastfed infant, or the effects of 3TC on milk production. Tenofovir Disoproxil Fumarate: Based on published data, tenofovir has been shown to be present in human breast milk (see Data). It is not known if

reactions in a breastfed infant similar to those seen in adults, instruct mothers not to breastfeed if they are receiving efavirenz, lamivudine and tenofovir disoproxil fumarate. Data: Tenofovir Disoproxil Fumarate: In a study of 50 HIV-uninfected, breastfeeding women on a tenofovir-containing regimen initiated between 1 and 24 weeks postpartum (median 13 weeks), tenofovir was undetectable in the plasma of most infants after 7 days of treatment in mothers. There were no serious adverse events in mothers or infants 8.3 Females and Males of Reproductive Potential

Because of potential teratogenic effects, pregnancy should be avoided in women receiving efavirenz, lamivudine and tenofovir disoproxil fumarate [see Warnings and Precautions (5.7), Use in Specific Populations (8.1)]. Pregnancy Testing: Females of reproductive potential should undergo pregnancy testing before initiation of efavirenz, lamivudine and tenofovir Contraception: Females of reproductive potential should use effective contraception during treatment with efavirenz, lamivudine and tenofovir

The safety and effectiveness of efavirenz, lamivudine and tenofovir disoproxil fumarate as a fixed-dose tablet in pediatric patients infected with HIV-1 and weighing at least 35 kg have been established based on clinical studies using the individual components (efavirenz, lamivudine, and tenofovir

enz, lamivudine and tenofovir disoproxil fumarate is not recommended for patients with impaired renal function (i.e., creatinine clearance less than 50 mL/min) or patients with end-stage renal disease (ESRD) requiring hemodialysis because it is a fixed-dose combination formulation that

If overdose occurs, the patient must be monitored for evidence of toxicity, and standard supportive treatment applied as necessary. Etavirenz: Some patients accidentally taking 600 mg twice daily have reported increased nervous system symptoms. One patient experienced Treatment of overdose with EFV should consist of general supportive measures, including monitoring of vital signs and observation of the patient's

dine and tenofovir disoproxil fumarate is a fixed-dose combination tablet for oral administration. Each film coated tablet contains efavirenz USP 400 mg, lamivudine USP 300 mg and tenofovir disoproxil fumarate 300 mg equivalent to tenofovir disoproxil 245 mg. Each tablet

Chloro-4-(cyclopropylethynyl)-1,4-dihydro-4-(trifluoromethyl)-2H-3,1-benzoxazin-2-one. Its molecular formula is C<sub>1</sub>,H<sub>2</sub>CIF<sub>2</sub>NO<sub>2</sub> and its structural

isopropoxycarbonyloxymethyl ester derivative of tenofovir. TDF is converted in vivo to tenofovir, an acyclic nucleoside phosphonate (nucleoside phosphonate) analog of adenosine 5'-monophosphate. Tenofovir exhibits activity against HIV-1 reverse transcriptase. The chemical name of TDF is  $9-[(R)-2-[[bis][(isopropoxycarbony])oxy]methoxy]phosphinyl]methoxy]propyl]adenine fumarate (1:1). It has a molecular formula of <math>C_{v_0}H_{v_0}N_{v_0}P^{-}C_{v_0}H_{v_0}O$ 

12 CLINICAL PHARMACOLOGY

Cardiac Electrophysiology: The effect of EFV on the QTc interval was evaluated in an open-label, positive and placebo-controlled, fixed single

relationship, the mean QTc prolongation and its upper bound 90% confidence interval are 8.7 ms and 11.3 ms in subjects with CYP2B6\*6/\*6

Efavirenz: EFV has been shown to pass into human breast milk. There is no information available on the effects of EFV on the breastfed infant, or the Lamivudine: 3TC is present in human milk. Samples of breast milk obtained from 20 mothers receiving 3TC monotherapy, 300 mg twice daily (2 times the dose in efavirenz, lamivudine and tenofovir disoproxil fumarate tablets), had measurable concentrations of 3TC. There is no information

Drug interactions with EFV are summarized in Table 5 [for pharmacokinetics data see Clinical Pharmacology (12.3, Tables 8 and 9)]. This table Table 5. Established and Other Potentially Significant Drug Interactions with EFV: Alteration in Dose or Regimen May Be Recommended tenofovir affects milk production or has effects on the breastfed child. Because of the potential for (1) HIV transmission (in HIV-negative infants): (2) developing viral resistance (in HIV-positive infants); and (3) adverse

disoproxil furmarate and for 12 weeks after discontinuing etavirenz, lamivudine and tenofovir disoproxil furmarate due to the long half-life of EFV. Barrier contraception should always be used in combination with other methods of contraception. Hormonal methods that contain progesterone may have decreased effectiveness [see Drug Interactions (7.5)].

disoproxil fumarate) 8.5 Geriatric Use Clinical studies of efavirenz, lamivudine and tenofovir disoproxil fumarate did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. In general, caution should be exercised in the administration of 3TC in elderly patients reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy. 8.6 Renal Impairment

cannot be adjusted [see Dosage and Administration (2.3)]. 8.7 Hepatic Impairment Efavirenz, lamivudine and tenofovir disoproxil fumarate is not recommended for patients with moderate or severe hepatic impairment because there are insufficient data to determine whether dose adjustment is necessary. Patients with mild hepatic impairment may be treated with efavirenz, udine and tenofovir disoproxil fumarate without any adjustment in dose [see Dosage and Administration (2.4), Warnings and Precautions (5.9) and Clinical Pharmacology (12.3)].

10 OVERDOSAGE

clinical status. Administration of activated charcoal may be used to aid removal of unabsorbed drug. There is no specific antidote for overdose with efavirenz. Since efavirenz is highly protein bound, dialysis is unlikely to significantly remove the drug from blood. Lamivudine: There is no known specific treatment for overdose with 3TC. If overdose occurs, the patient should be monitored and standard supportive treatment applied as required because a negligible amount of 3TC was removed via (4-hour) hemodialysis, continuous ambulatory peritoneal dialysis, and automated peritoneal dialysis, it is not known if continuous hemodialysis would provide clinical benefit in a 3TC overdose Tenofovir Disoproxil Fumarate: Limited clinical experience at doses higher than the therapeutic dose of TDF 300 mg is available. Tenofovir is efficiently removed by hemodialysis with an extraction coefficient of approximately 54%. Following a single 300 mg dose of tenofovir disoproxil fumarate, a 4-hour hemodialysis session removed approximately 10% of the administered tenofovir dose.

11 DESCRIPTION contains the following inactive ingredients: croscarmellose sodium, ferric oxide yellow, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium lauryl sulfate, talc and titanium dioxide. Efavirenz: Efavirenz (EFV) is an HIV-1 specific, non-nucleoside, reverse transcriptase inhibitor (NNRTI). Efavirenz is chemically described as (S)-6-

Efavirenz USP is a white to off white powder with a molecular mass of 315.67. It is soluble in methanol and practically insoluble in water (< 10

Lamivudine: Lamivudine (also known as 3TC) is a synthetic nucleoside analogue with activity against HIV-1 and HBV. The chemical name of lamivudine is (-)-1-1([R,5]-2-(Hydroxy methyl)-1,3-oxathiolan-5-yl]cytosine. Lamivudine is the (-)enantiomer of a dideoxy analogue of cytidine. Lamivudine has also been referred to as (-)2',3' -dideoxy, 3' -thiacytidine. It has a molecular formula of  $C_8H_{11}N_3O_3S$  and a molecular weight of 229.26 g per mol. It has the following structural form

Lamivudine USP is a white or almost white powder and is soluble in water. Tenofovir Disoproxil Fumarate: Tenofovir disoproxil fumarate (TDF) (a prodrug of tenofovir) is a fumaric acid salt of bis-

Tenofovir disoproxil fumarate is a white to off white powder freely soluble in dimethyl formamide, soluble in methanol and slightly soluble in water at 25°C. It has a partition coefficient (log p) of 0.75.

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are a fixed-dose combination of antiviral drugs EFV, 3TC, and TDF with antiviral activity against HIV-1 [see Microbiology (12.4)].

microgram/mL).

sequence 3-period, 3-treatment crossover QT study in 58 healthy subjects enriched for CYP2B6 polymorphisms. The mean C<sub>max</sub> of EFV in subjects with CYP2B6 \*6/\*6 genotype following the administration of 600 mg daily dose for 14 days was 2.25-fold the mean C<sub>max</sub> observed in subjects with CYP2B6 \*1/\*1 genotype. A positive relationship between EFV concentration and QTc prolongation was observed. Based on the concentration-QTc

irenz, lamivudine and **Call your healthcare** 

think you may be pregnant or become pregnant during treatment with efavirenz, lamivudine and who are able to become pregnant should use effective birth control during treatment with efavirenz, lamivudine and disoproxil fumarate tablets and for 12 weeks after stopping treatment. A barrier form of birth control should always be used able to become pregnant, your healthcare provider should do a pregnancy test before you start efavirenz, lamivudine ar disoproxil fumarate tablets. lamivudine and tenofovir health of you and your ba

No dosage adjustment is recommended when efavirenz, lamivudine and tenofovir disoproxil fumarate is administered with the following: aluminum/magnesium hydroxide antacids, azithromycin, cetirizine, famotidine, fluconazole, and lorazepam 7.7 Drugs Inhibiting Organic Cation Transporters 3TC, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, is predominantly eliminated in the urine by active organic cationic secretion. The possibility of interactions with other drugs administered concurrently should be considered, particularly when their main route of elimination is active renal secretion via the organic cationic transport system (e.g., trimethoprim) [see Clinical Pharmacology (12.3)]. No data are available regarding interactions with other drugs that have renal clearance mechanisms similar to that of 3TC.

6% Vomiting 8% Arthralgia  $Frequencies \ of adverse\ reactions\ are\ based\ on\ all\ treatment-emergent\ adverse\ events, regardless\ of\ relationship\ to\ study\ drug.$ Bash event includes rash, pruritus, maculopapular rash, urticaria, vesiculobullous rash, and pustular rash, Lipodystrophy represents a variety of investigator-described adverse events not a protocol-defined syndrom Peripheral neuropathy includes peripheral neuritis and neuropathy.

ENCORE1 Study - Adverse Reactions: The most common adverse reactions seen in a double-blind comparative controlled study in which 630 reatment-naïve subjects received EFV 400 mg (N = 321) or EFV 600 mg (N = 309) in combination with fixed-dose emtricitabine (FTC)/TDF for 48 weeks were mild to moderate gastrointestinal events, dizziness, abnormal dreams, and rash. Selected clinical adverse reactions of moderate o

	EFV 400 mg + FTC/TDF	EFV 600 mg + FTC/TDI
	N = 321	N = 309
ash event <sup>b</sup>	9%	13%
zziness	6%	9%
somnia	3%	4%
onormal dreams	2%	2%
eadache	1%	3%
arrhea	2%	3%
omiting	1%	2%
rexia	2%	1%
oper respiratory tract infection	3%	1%
asopharyngitis	3%	2%
erpes zoster	3%	1%
stroenteritis	2%	2%

generalized, rash macular, rash maculopapular, rash morbilliform, rash papular, rash pruritic, rash vesicular, and urticaria. Laboratory Abnormalities: Table 3 provides a list of laboratory abnormalities (Grades 3 to 4) observed in Trial 903. With the exception of fasting cholesterol and fasting triglyceride elevations that were more common in the d4T group (40% and 9%) compared with the TDF group (19% and 1%) respectively, laboratory abnormalities observed in this trial occurred with similar frequency in the TDF and d4T treatment arms. Table 3. Grade 3 to 4 Laboratory Abnormalities Reported in ≥1% of Patients Randomized to Efavirenz, Lamivudine and Tenofovir Disoproxi Fumarate in Study 903 (0 to 144 Weeks)

	TDF + 3TC + EFV	d4T + 3TC + EFV
	N = 299	N = 301
Any ≥ Grade 3 Laboratory Abnormality	36%	42%
Fasting Cholesterol (> 240 mg/dL)	19%	40%
Creatine Kinase (M: > 990 U/L; F: > 845 U/L)	12%	12%
Serum Amylase (> 175 U/L)	9%	8%
AST (M: > 180 U/L; F: > 170 U/L)	5%	7%
ALT (M: > 215 U/L; F: > 170 U/L)	4%	5%
Hematuria (> 100 RBC/HPF)	7%	7%
Neutrophils (< 750/mm³)	3%	1%
Fasting Triglycerides (> 750 mg/dL)	1%	9%
ENCORE1 study, a summary of Grade 3 and 4 laboratory abn ole 4. Grades 3 to 4 Laboratory Abnormalities in ≥ 2% in Ei		
Lahoratory Parameter		

Laboratory Parameter	EFV 400 mg + FTC + TDF	EFV 600 mg + FTC + TDF
	N = 321	N = 309
	5%	3%
	2%	2%

and may be associated with increased efavirenz exposure. Patients with serious psychiatric adverse experiences should seek immediate medical

evaluation to assess the possibility that the symptoms may be related to the use of EFV, and if so, to determine whether the risks of continued

There have also been occasional postmarketing reports of death by suicide, delusions, psychosis-like behavior, although a causal relationship to the use of FFV cannot be determined from these reports [see Adverse Reactions (6.2)]. Postmarketing cases of catatonia have also been reported tenofovir

of the most common (occurring in > 1% patients) psychiatric events among patients who received EFV 400 mg (N = 321) or EFV 600 mg (N = 309) regimens, respectively, were: abnormal dreams (8.7%, 11.3%), insomnia (6.2%, 6.5%), somnolence (3.1%, 3.9%), depression (3.1%, 1.6%), nightmare (1.9%, 2.6%), sleep disorder (2.2%, 1.3%), and anxiety (1.2%, 1.3%).

istration of single doses of 3TC and sorbitol resulted in a sorbitol dose-dependent reduction in 3TC exposures. When possible, avoid use of sorbitol-containing medicines with 3TC [see Clinical Pharmacology (12.3)].

8 USE IN SPECIFIC POPULATIONS 8.1 Pregnancy lamivudine and tenofovir disoproxil fumarate during pregnancy. Healthcare providers are encouraged to register patients by calling the ı get a new to

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vita herbal supplements.
Some medicines interact with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Efavirenz, lamivudine and tenofovir cfumarate tablets may affect the way other medicines work, and other medicines may affect how efavirenz, lamivudine and disoproxil fumarate tablets works. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you medicine. healthcare provider or pharmacist for a list of medicines that interact with efavirenz,

disoproxil fumarate tablets. Do not start taking a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets with other medicines. efavirenz, lamivudine and tenorovir disoproxil fumarate tablets?
Take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets exactly as your healthcare provider tells you to take it.
Take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets 1 time each day, preferably at bedtime. Taking efavirenz,
Take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets 5 time each day, preferably at bedtime. Taking efavirenz,

and tenofovir disoproxil fumarate tablets at bedtime might help to make some of the side effects less bothersome.

Take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets on an empty stomach.

Do not miss a dose of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. If you miss a dose, take the missed dose as soon as you remember. If it is almost time for your next dose of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, do not take the

become harder to treat. When your supply starts to run low, get more from your healthcare provider or pharmacy. If you take too much efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, go to the nearest hospital Do not miss a dose of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. If you miss as you remember. If it is almost time for your next dose of efavirenz, lamivudine and tenofovir dison missed dose. Take the next dose at your regular time.
Stay under the care of your healthcare provider during treatment with efavirenz, lamivudine and ter Do not run out of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. The virus in your

increase and the virus

ofovir disoproxil fum blood may increase a

What should I avoid while taking efavirenz, lamivudine and tenofovir disoproxil fumarate tablets?
You should avoid taking medicines that contain sorbitol during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. What are the possible side effects of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets?

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets may cause serious side effects, including:
• See "What is the most important information I should know about efavirenz, lamivudine and tenofovir disoproxil fumarate tablets?" tenofovir disoproxil fumarate tablets. Lactic acidosis is a serious medical emergency that can le **provider right away if you get any of the following symptoms that could be signs of lactic acidosis:**o feel very weak or tired

feel very weak or tired **Build-up of an acid in your blood (lactic acidosis).** Lactic acidosis tenofovir disoproxil fumarate tablets. Lactic acidosis is a serious m

Severe liver problems can happen in people who take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. In some cases, these severe liver problems can lead to death. Your liver may become large (hepatomegaly) and you may develop fat in your liver (steatosis). Inflammation of your liver (hepatitis) that can lead to liver failure requiring a liver transplant has been reported in some people treated with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Your healthcare provider may do blood tests to check your liver before and during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. feel dizzy or lighthea have a fast or irregul unusual (not normal) muscle pain trouble breathing

**Call your healthcare provider right away if you get any of the following signs or symptoms of liver problems:**your skin or the white part of your eyes turns
loss of appetite for several days or longer , aching, or tendernes of your stomach-area 0 0 0 0 0

500 mm

Leaflet Size: 830X500 MM Font: Helvetica Condensed Font Size: 6 pt Colour: Black Pharma Code: 679

HIV-1 is the virus that causes AIDS (Acquired Immune Deficiency Syndrome).
Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets contains the prescription medicines efavirenz, disoproxil fumarate.
Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are not for use in children weighing less than 35 kg.

Before you take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, tell your healthcare provider about all of your conditions, including if you:

Do not take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets if you:
 are allergic to efavirenz, lamivudine, tenofovir disoproxil fumarate, or any of the ingredients in efavirenz, lamivudine and i disoproxil fumarate tablets. See the end of this Patient Information leaflet for a complete list of ingredients in efavirenz, lamivu tenofovir disoproxil fumarate tablets.

are currently taking elbasvir and grazoprevir.

have liver problems, including hepatitis B or C infection have kidney problems, including end-stage renal disease (ESRD) that requires dialysis

have a history of mental health problems
have a history of drug or alcohol abuse
have a heart problem, including QT prolongation
have bone problems, including a history of bone fractures
have a history of seizures
are pregnant or plan to become pregnant. Efavirenz, lamivudine and tenofovir disoproxil or you should not become pregnant during treatment with efavirenz, lamivudine and healthcare provider right away if you think you may be pregnant or become pregnant tenofovir disoproxil furnarate tablets.

along with another type If you are able to becon

**Pregnancy Registry.** There is a pregnancy registry for women who take efavirenz, during pregnancy. The purpose of this registry is to collect information about the provider about how you can take part in this registry.

breastfeeding or plan to breastfeed. Do not breastfeed if you take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. Talk to your healthcare provider about the best way to feed your baby.

than 50 mL/min or patients with end-stage renal disease requiring hemodialysis. (2.3) Hepatic Impairment: Not recommended for patients with moderate or severe hepatic impairment. Use caution in patients with mild hepatic impairment. (2.4) ----DOSAGE FORMS AND STRENGTHS----Tablets: 400 mg efavirenz, 300 mg lamivudine and 300 mg tenofovir

stomach, preferably at bedtime. (2.2)

patients weighing at least 35 kg. (1)

obtained. (2.1)

HIGHLIGHTS OF PRESCRIBING INFORMATION

TABLETS.

ets, for oral use

Initial U.S. Approval: 2018

anti-hepatitis B treatment. (5.1)

Warnings and Precautions, Use with Interferon- and Ribavirin-Based Regimens (previous 5.10)

These highlights do not include all the information needed to use

EFAVIRENZ, LAMIVUDINE AND TENOFOVIR DISOPROXIL FUMARATE

TABLETS safely and effectively. See full prescribing information for EFAVIRENZ, LAMIVUDINE AND TENOFOVIR DISOPROXIL FUMARATE

EFAVIRENZ. LAMIVUDINE AND TENOFOVIR DISOPROXIL FUMARATE

WARNING: POST TREATMENT ACUTE EXACERBATIONS OF

HEPATITIS B

 ${\it See full prescribing information for complete boxed warning.}$ 

Severe acute exacerbations of hepatitis B have been reported

in patients who are co-infected with HBV and human

lamivudine and tenofovir disoproxil fumarate. Monitor hepatic

function closely in these patients and, if appropriate, initiate

Warnings and Precautions (5.1, 5.2, 5.3, 5.4, 5.6, 5.13, 5.14) 10/2019

immunodeficiency virus (HIV-1) and have discontinu

-----RECENT MAJOR CHANGES---

----INDICATIONS AND USAGE---

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are three-drug combination of efavirenz (EFV), a non-nucleoside reverse

transcriptase inhibitor, and lamivudine (3TC) and tenofovir disoproxil fumarate (TDF), both nucleo(t)side reverse transcriptase inhibitors and

are indicated as a complete regimen for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adult and pediatric

· Testing: Prior to initiation and during treatment with efavirenz,

Renal Impairment: Not recommended in patients with CrCL less

should be tested for hepatitis B virus infection, and estimated

creatinine clearance, urine glucose, and urine protein should be

Recommended dose: One tablet taken orally once daily on an empty
 FDA-1088 or www.fda.gov/medwatch.

----DOSAGE AND ADMINISTRATION----

Serious Psychiatric Symptoms: Immediate medical evaluation is

recommended for serious psychiatric symptoms such as severe depression or suicidal ideation. (5.5)

Nervous System Symptoms (NSS): NSS are frequent, usually begin

1 to 2 days after initiating therapy and resolve in 2 to 4 weeks.

of onset of psychiatric symptoms. (5.6)

pre-existing hepatic disease. (5.9, 8.7)

fumarate as clinically appropriate. (5.10)

therapy and periodically thereafter. (5.12)

develops. (5.8)

or bone loss. (5.13)

udine and tenofovir disoproxil fumarate tablets, patients • Most common adverse reactions (>5% with efavirenz, lamivudine

Dosing at bedtime may improve tolerability. NSS are not predictive  $% \left( 1\right) =\left( 1\right) \left( 1\right)$ 

Rash: Rash usually begins within 1 to 2 weeks after initiating

therapy and resolves within 4 weeks. Discontinue if severe rash

Hepatotoxicity: Monitor liver function tests before and during

treatment in patients with underlying hepatic disease, including

hepatitis B or C coinfection, marked transaminase elevations, or

who are taking medications associated with liver toxicity. Among

reported cases of hepatic failure, a few occurred in patients with no

Pancreatitis: Use with caution in pediatric patients with a history of

pancreatitis or other significant risk factors for pancreatitis.

Discontinue efavirenz, lamivudine and tenofovir disoproxil

Convulsions: Use caution in patients with a history of seizures.

Lipids: Total cholesterol and triglyceride elevations. Monitor before

Decreases in Bone Mineral Density (BMD): Observed in HIV-

infected patients. Consider assessment of BMD in patients with a

history of pathologic fracture or other risk factors for osteoporosis

Immune Reconstitution Syndrome: Observed in HIV-infected

patients. May necessitate further evaluation and treatment. (5.14)

Redistribution/Accumulation of Body Fat: Observed in HIV-infected

patients receiving antiretroviral combination therapy. (5.15)

-----ADVERSE REACTIONS-----

and tenofovir disoproxil fumarate) are rash and dizziness. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Laurus

-----DRUG INTERACTIONS-

treatment of HIV-1 infection (7.1)

transmission. (8.2)

7 DRUG INTERACTIONS

7.8 Sorbitol

8.1 Pregnancy 8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

8.6 Renal Impairment

8.7 Hepatic Impairment

12 CLINICAL PHARMACOLOGY

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOG

12.4 Microbiology

14 CLINICAL STUDIES

are not listed.

Tablets: 400 mg of efavirenz, 300 mg of lamivudine, and 300 mg of tenofovir disoproxil fumarate (equivalent to 245 mg of tenofovir disoproxil).

Important Differences Among Lamivudine-Containing Products: Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets contain a higher

Loss of therapeutic effect of efavirenz, lamivudine and tenofovir disoproxil fumarate and possible development of resistance.

Possible clinically significant adverse reactions from greater exposures of concomitant drugs

12.1 Mechanism of Action

7.2 QT Prolonging Drugs

8 USE IN SPECIFIC POPULATIONS

7.3 Drugs Affecting Renal Function7.4 Cannabinoid Test Interaction

patient labeling.

Generics Inc. at 1-833-3-LAURUS (1-833-352-8787) or FDA at 1-800-

Efavirenz lamivudine and tenofovir disoproxil fumarate should not

be administered with other antiretroviral medications for the

Coadministration of efavirenz, lamivudine and tenofovir disoproxil

fumarate can alter the concentrations of other drugs and other

drugs may alter the concentration of efavirenz, lamivudine and

tenofovir disoproxil fumarate. The potential for drug-drug

interactions should be considered before and during therapy. (5.3.

Pregnancy: Women should avoid pregnancy during EFV therapy, a

component of efavirenz, lamivudine and tenofovir disoproxi

Lactation: Breastfeeding not recommended due to potential for HIV

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved

7.1 Not Recommended with Other Antiretroviral Medications

7.5 Established and Other Potentially Significant Interactions
 7.6 Drugs without Clinically Significant Interactions

7.7 Drugs Inhibiting Organic Cation Transporters

8.3 Females and Males of Reproductive Potential

13.1 Carcinogenesis. Mutagenesis. Impairment of Fertility

Sections or subsections omitted from the full prescribing information

13.2 Animal Toxicology and/or Pharmacology

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

14.1 Clinical Efficacy in Patients with HIV-1 Infection

and contraception are recommended. (8.3)

fumarate tablets, and for 12 weeks after discontinuation. (5.7, 8.1,

Females and Males of Reproductive Potential: Pregnancy testing

--- USE IN SPECIFIC POPULATIONS---

disoproxil fumarate (equivalent to 245 mg of tenofovir disoproxil). (3) -----CONTRAINDICATIONS-----. Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are contraindicated in patients with previous hypersensitivity (e.g., Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of this product. (4) Coadministration with elbasvir/grazoprevir. (4) ----WARNINGS AND PRECAUTIONS----

 Lactic Acidosis/Severe Hepatomegaly with Steatosis: Discontinue treatment in patients who develop symptoms or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity. (5.2) New Onset or Worsening Renal Impairment: Can include acute renal failure and Fanconi syndrome. Avoid administering efavirenz, lamivudine and tenofovir disoproxil fumarate tablets with concurrent or recent use of nephrotoxic drugs. (5.4) FULL PRESCRIBING INFORMATION: CONTENTS\*

WARNING: POST TREATMENT ACUTE EXACERBATIONS OF HEPATITIS B 1 INDICATIONS AND USAGE 2 DOSAGE AND ADMINISTRATION 2.1 Testing Prior to Initiation and During Treatment with Efavirenz, Lamiyudine and Tenofovir Disoproxil Fumarate Tablets 2.2 Recommended Dosage for Adult and Pediatric Patients Weighing at Least 35 kg 2.3 Not Recommended in Renal Impai 2.4 Not Recommended in Moderate to Severe Hepatic Impairment 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS

5.1 Severe Acute Exacerbation of Hepatitis B in Patients Coinfected with HIV-1 and HBV to Drug Interactions 5.4 New Onset or Worsening Renal Impairment 5.5 Psychiatric Symptoms 5.6 Nervous System Symptoms 5.7 Embryo-Fetal Toxicity 5.10 Pancreatitis 5.12 Lipid Elevations

5.2 Lactic Acidosis and Severe Hepatomegaly with Steatosis 5.3 Risk of Adverse Reactions or Loss of Virologic Response Due 5.8 Skin and Systemic Hypersensitivity Reaction 5.13 Bone Loss and Mineralization Effects

5.14 Immune Reconstitution Syndrome 5.16 QTc Prolongation

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

FULL PRESCRIBING INFORMATION

6 ADVERSE REACTIONS

WARNING: POST TREATMENT ACUTE EXACERBATIONS OF HEPATITIS B

Severe acute exacerbations of hepatitis B have been reported in patients who are co-infected with hepatitis B virus (HBV) and human immunodeficiency virus (HIV-1) and have discontinued lamivudine or tenofovir disoproxil fumarate, two components of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Monitor benatic function closely in these natients and, if appropriate, initiate antihepatitis B treatment [see Warnings and Precautions (5.1)] I INDICATIONS AND USAGE enz, lamivudine and tenofovir disoproxil fumarate tablets are indicated as a complete regimen for the treatment of human immunodeficiency

virus type 1 (HIV-1) infection in adult and pediatric patients weighing at least 35 kg. 2 DOSAGE AND ADMINISTRATION 2.1 Testing Prior to Initiation and During Treatment with Efavirenz, Lamiyudine and Tenofovir Disoproxil Fumarate Tablets Prior to initiation of efavirenz, lamivudine and tenofovir disoproxil furnarate tablets, test patients for hepatitis B virus infection [see Warnings and Precautions (5.1)1. It is recommended that serum creatinine, serum phosphorus, estimated creatinine clearance, urine glucose, and urine protein be assessed before initiating efavirenz, lamivudine and tenofovir disoproxil fumarate tablets and during therapy in all patients as clinically appropriate [see Warnings and Precautions (5.4)1. Monitor hepatic function prior to and during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets [see Warnings and Precautions (5.9)]. 2.2 Recommended Dosage for Adult and Pediatric Patients Weighing at Least 35 kg

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are a three-drug fixed-dose combination product containing 400 mg of efavirenz (EFV), 300 mg of lamivudine (3TC), and 300 mg of tenofovir disoproxil fumarate (TDF). The recommended dosage of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets in HIV-1-infected adults and pediatric patients weighing at least 35 kg is one tablet taken orally once daily. Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets should be taken on an empty stomach, preferably at bedtime. Dosing at bedtime may improve the tolerability of nervous system symptoms [see Warnings and Precautions (5.6) and Adverse Reactions (6.1)]. 2.3 Not Recommended in Renal Impairment Because efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are a fixed-dose combination tablet and cannot be dose adjusted, they are not recommended for patients with impaired renal function (creatinine clearance less than 50 mL/min) or patients with end-stage renal disease (ESRD) requiring hemodialysis [see Use in Specific Populations (8.6)].  ${\bf 2.4}\quad {\bf Not\,Recommended\,in\,Moderate\,to\,Severe\,Hepatic\,Impairment}$ Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are not recommended in patients with moderate or severe hepatic impairment (Child-Pugh B or C) [see Warnings and Precautions (5.9) and Use in Specific Populations (8.7)].

The 400 mg/300 mg/300 mg tablets are white to off-white, film-coated, oval, biconvex tablets debossed with "L40" on one side and plain on other 4 CONTRAINDICATIONS Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are contraindicated: • in patients with a previous hypersensitivity reaction (e.g., Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components contained in the formulation [see Warnings and Precautions (5.8)]. when coadministered with elbasvir and grazoprevir [see Warnings and Precautions (5.3) and Drug Interactions (7.5)]. WARNINGS AND PRECAUTIONS 5.1 Severe Acute Exacerbation of Hepatitis B in Patients Coinfected with HIV-1 and HBV Posttreatment Exacerbations of Hepatitis: All patients with HIV-1 should be tested for the presence of chronic hepatitis B virus (HBV) before initiating antiretroviral therapy *[see Dosage and Administration (2.1)]*. Discontinuation of anti-HBV therapy, including 3TC and TDF, may be associated with severe acute exacerbations of hepatitis B. Patients infected with HBV who discontinue efavirenz, lamivudine and tenofor disoproxil fumarate should be closely monitored with both clinical and laboratory follow-up for at least several months after stopping treatment. If appropriate, resumption of anti-hepatitis B therapy may be warranted.

3 DOSAGE FORMS AND STRENGTHS

dose of the same active ingredient, 3TC, than EPIVIR-HBV® tablets. EPIVIR-HBV was developed for patients with chronic hepatitis B. The formulation and dosage of 3TC in EPIVIR-HBV are not appropriate for patients co-infected with HIV-1 and HBV. Safety and efficacy of 3TC have not been established for treatment of chronic hepatitis B in patients co-infected with HIV-1 and HBV. If treatment with EPIVIR-HBV, TDF, or a tenofovir alafenamide (TAF)-containing product is prescribed for chronic hepatitis B for a patient with unrecognized or untreated HIV-1 infection, rapid emergence of HIV-1 resistance is likely to result because of the subtherapeutic dose and the inappropriateness of monotherapy HIV-1 treatment. 5.2 Lactic Acidosis and Severe Hepatomegaly with Steatosis Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogs and other antiretrovirals. Female sex and obesity may be risk factors for the development of lactic acidosis and severe hepatomegaly with steatosis in patients treated with antiretroviral nucleoside analogues. Treatment should be suspended in any patient who develops clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity, which may include hepatomegaly and steatosis even in the absence of marked 5.3 Risk of Adverse Reactions or Loss of Virologic Response Due to Drug Interactions The concomitant use of efavirenz, lamivudine and tenofovir disoproxil fumarate and other drugs may result in known or potentially significant drug interactions, some of which may lead to [see Contraindications (4) and Drug Interactions (7.5)]:

See Table 5 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations. Consider the potential for drug interactions prior to and during therapy with efavirenz, lamivudine and tenofovir disoproxil fumarate; review concomitant medications during therapy with efavirenz, lamivudine and tenofovir disoproxil fumarate; and monitor for the adverse reactions associated with the concomitant drugs. 5.4 New Onset or Worsening Renal Impairment TDF, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets is principally eliminated by the kidney. Renal impairment including cases of acute renal failure and Fanconi syndrome (renal tubular injury with severe hypophosphatemia), has been reported with the use of TDF I see Adverse Reactions (6.2)1 Prior to initiation and during use of efavirenz, lamivudine and tenofovir disoproxil fumarate, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, urine glucose, and urine protein in all patients. Avoid efavirenz, lamivudine and tenofovir disoproxil fumarate with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple nonsteroidal anti-inflammatory drugs [NSAIDs]) (see Drug Interactions (7.3)). Cases of acute renal failure after initiation of high-dose or multiple NSAIDs have been reported in HIV-infected patients with risk factors for renal dysfunction who appeared stable on TDF. Some patients required hospitalization and renal replacement therapy. Alternatives to NSAIDs should be considered, if needed, in patients at risk for renal dysfunction. Persistent or worsening bone pain, pain in extremities, fractures and/or muscular pain or weakness may be manifestations of proximal renal

ppathy and should prompt an evaluation of renal function in patients at risk of renal dysfunction 5.5 Psychiatric Symptoms ous psychiatric adverse experiences have been reported in patients treated with EFV, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. In controlled trials of 1,008 patients treated with regimens containing EFV for a mean of 2.1 years and 635 patients treated with control regimens for a mean of 1.5 years, the frequency (regardless of causality) of specific serious psychiatric events among patients who received EFV or control regimens, respectively, were severe depression (2.4%, 0.9%), suicidal ideation (0.7%, 0.3%), nonfatal suicide attempts (0.5%, 0), aggressive behavior (0.4%, 0.5%), paranoid reactions (0.4%, 0.3%), and manic reactions (0.2%, 0.3%). When psychiatric symptoms similar to those noted above were combined and evaluated as a group in a multifactorial analysis of data from a study using EFV 600 mg, treatment with EFV was associated with an increase in the occurrence of these selected psychiatric symptoms. Other factors associated with an increase in the occurrence of these psychiatric symptoms were history of injection drug use, psychiatric history, and receipt of psychiatric medication at study entry; similar associations were observed in both the EFV and control treatment groups. In a study using EFV 600 mg, onset of new serious psychiatric symptoms occurred throughout the study for both EFV-treated and control-treated patients. One percent of EFV-treated patients discontinued or interrupted treatment because of one or more of these selected psychiatric symptoms. In the ENCORE1 (Evaluation of Novel Concepts in Ontimization of antiRetroviral Efficacy) study at Week 48 the frequency (regardless of causality)

Patient Information
Efavirenz, Lamivudine and Tenofovir Disoproxil Fumarate
(ef-a-vir-enz, Ia-MIV-ue-deen and ten-OF-oh-vir DYE-soe-PROX-il FUE-ma-rate)

dark or "tea-colored" urine light-colored stools (bowel movements)

шш

Whatare efavirenz, lamivudine and tenofovir disoproxil fumarate tablets? Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets is a prescription medicine that is used without other antiretroviral medicines to treat Human Immunodeficiency Virus-1 (HIV-1) in people weighing at least 35 kg.

more information about side effects, see "What are the possible side effects of efavirenz, lamivudine and tenofovir disoproxil arate tablets?"

most important information I should know about efavirenz, Iamivudine and tenofovir disoproxil fumarate tablets? amivudine and tenofovir disoproxil fumarate tablets can cause serious side effects, including:

provider before your efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are all gone.

Do not stop efavirenz, lamivudine and tenofovir disoproxil fumarate tablets without first talking to your healthcare provider. If you stop taking efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, your healthcare provider will need to check your health often and do blood tests regularly for several months to check your liver. **Worsening of Hepatitis B virus infection.** If you have Human Immunodeficiency Virus type 1 (HIV-1) and Hepatitis B Virus (HBV) infection, your HBV may get worse (flare-up) if you stop taking efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. A "flare-up" is when your HBV infection suddenly returns in a worse way than before. Your healthcare provider will test you for HBV infection before you start treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. up" is when your HBV infection suddenly returns in a worse way than before. Your healthcare provider will test you for HBV infe before you start treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are safe and effective in people who have both land HBV infection.

Do not run out of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Refill your prescription or talk to your healt

The effect of food on efavirenz, lamivudine and tenofovir disoproxil fumarate has not been evaluated. Efavirenz: In HIV-1-infected subjects, time-to-peak plasma concentrations were approximately 3 to 5 hours and steady-state plasma concentrations were reached in 6 to 10 days. EFV is highly bound (approximately 99.5 to 99.75%) to human plasma proteins, predominantly albumin. Following administration of 14C-labeled EFV, 14 to 34% of the dose was recovered in the urine (mostly as metabolites) and 16 to 61% was recovered in feces (mostly as parent drug). *In vitro* studies suggest CYP3A and CYP2B6 are the major isozymes responsible for EFV metabo EFV has been shown to induce CYP enzymes, resulting in induction of its own metabolism. EFV has a terminal half-life of 52 to 76 hours after single

doses and 40 to 55 hours after multiple doses. Lamivudine: After oral administration of 2 mg/kg of 3TC twice a day to 9 adults with HIV-1, the peak serum 3TC concentration (C<sub>mv</sub>) was 1.5 ± 0.5 mcg/mL (mean ± SD). The area under the plasma concentration versus time curve (AUC) and C<sub>max</sub> increased in proportion to oral dose over the range from 0.25 to 10 mg/kg and absolute bioavailability in 12 adult patients was 86% ± 16% (mean ± SD) for the 150-mg tablet and 87% ± 13% for the oral solution. Binding of 3TC to human plasma proteins is low (< 36%). Within 12 hours after a single oral dose of 3TC in 6 HIV-1-infected adults, 5.2% ± 1.4% (mean ± SD) of the dose was excreted as the trans-sulfoxide metabolite in the urine. The majority of 3TC is eliminated unchanged in urine by active organic cationic secretion and the observed mean elimination half-life (t<sub>1/2</sub>) ranged from 5 to 7 hours in most single-

dose studies with serum sampling for 24 hours after dosing. **Tenafovir Disoproxil Fumarate:** Following oral administration of a single 300 mg dose of TDF to HIV-1-infected subjects in the fasted state, maximum serum concentrations ( $C_{max}$ ) were achieved in 1.0  $\pm$  0.4 hrs (mean  $\pm$  SD) and  $C_{max}$  and AUC values were 296  $\pm$  90 ng/mL and 2,287  $\pm$  685 ng-hr/mL, respectively. The oral bioavailability of tenofovir from TDF in fasted subjects is approximately 25%. Less than 0.7% of tenofovir binds to human plasma proteins *in vitro* and the binding is independent of concentration over the range of 0.01 to 25 mcg/mL. Approximately 70 to 80% of the intravenous dose of tenofovir is recovered as unchanged drug in the urine. Tenofovir is eliminated by a combination of glomerular filtration and active tubular secretion with a renal clearance in adults with normal renal function of  $243 \pm 33$  mL/min (mean  $\pm$  SD). Following a single oral dose, the  $terminal\ elimination\ half-life\ of\ tenofovir\ is\ approximately\ 17\ hours.$ 

Special Populations:

Efavirenz and Lamivudine: There are no significant or clinically relevant racial differences in EFV and 3TC pharmacokinetics. Tenofovir Disoproxil Fumarate: There were insufficient numbers from racial and ethnic groups other than Caucasian to adequately determine potential pharmacokinetic differences among these populations Gender: There are no significant or clinically relevant gender differences in the pharmacokinetics of EFV, 3TC, and TDF.

Table 6. Pharmacokinetic Parameters (Mean ± SD) after a Single 300-mg Oral Dose of 3TC in Subjects with Varying Degrees of Renal Function

Geriatric Patients: The pharmacokinetics of 3TC and TDF have not been studied in patients over 65 years of age. Patients with Renal Impairment: [See Use in Specific Populations (8.6).]

Efavirenz: The pharmacokinetics of EFV have not been studied in patients with renal impairment <u>Lamivudine:</u> The pharmacokinetics of 3TC are altered in subjects with renal impairment (Table 6)

	Creatinine Clearance Criterion (Number of Subjects)					
Parameter	> 60 mL/min (n = 6)	10 to 30 mL/min (n = 4)	< 10 mL/mii (n = 6)			
Creatinine clearance (mL/min)	111 ± 14	28 ± 8	6 ± 2			
C <sub>max</sub> (mcg/mL)	2.6 ± 0.5	3.6 ± 0.8	5.8 ± 1.2			
AUC <sub>∞</sub> (mcg•h/mL)	11.0 ± 1.7	48.0 ± 19	157 ± 74			
CI/F (mL/min)	464 ± 76	114 ± 34	36 ± 11			

Tenofovir Disoproxil Furnarate: The pharmacokinetics of TDF are altered in subjects with renal impairment [see Warnings and Precautions (5.4)]. In subjects with creatinine clearance below 50 mL/min or with end-stage renal disease (ESRD) requiring dialysis, Cmear and AUC, and AUC the of tenofovir were

ncreased.		,	, , , , , , , , , , , , , , , , , , , ,	, 0000	
able 7. Pharmacokinetic Pa enal Function	arameters (Mean ± SD) of T	enofovir After a Single 300-r	ng Oral Dose of TDF in Subj	ects with Varying Degrees	s of
Baseline Creatinine Clearance (mL/min)	> 80 (N = 3)	50 to 80 (N = 10)	30 to 49 (N = 8)	12 to 29 (N = 11)	
C <sub>max</sub> (mcg/mL)	0.34 ± 0.03	0.33 ± 0.06	0.37 ± 0.16	0.60 ± 0.19	٦
AUC <sub>□ to∞</sub> (mcg•hr/mL)	2.18 ± 0.26	3.06 ± 0.93	6.01 ± 2.50	15.98 ± 7.22	٦
CL/F (mL/min)	1,043.7 ± 115.4	807.7 ± 279.2	444.4 ± 209.8	177.0 ± 97.1	٦
CL <sub>renal</sub> (mL/min)	243.5 ± 33.3	168.6 ± 27.5	100.6 ± 27.5	43.0 ± 31.2	٦

Efavirenz: A multiple-dose study showed no significant effect on EFV pharmacokinetics in patients with mild hepatic impairment (Child-Pugh Class A) compared with controls. There were insufficient data to determine whether moderate or severe hepatic impairment (Child-Pugh Class B or C) affects EFV pharmacokinetics. Lamivudine: The pharmacokinetic properties of 3TC have been determined in adults with impaired hepatic function. Pharmacokinetic parameters were not altered by diminishing hepatic function. Safety and efficacy of 3TC have not been established in the presence of decompensated liver Tenofovir Disoproxil Fumarate: The pharmacokinetics of tenofovir following a 300 mg single dose of TDF have been studied in non-HIV infected subjects with moderate to severe (Child-Pugh B to C) hepatic impairment. There were no substantial alterations in tenofovir pharmacokinetics in

Assessment of Drug Interactions: [See Drug Interactions (7).] Efavirenz: EFV has been shown in vivo to cause hepatic enzyme induction, thus increasing the biotransformation of some drugs metabolized by CYP3A and CYP2B6. In vitro studies have shown that EFV inhibited CYP isozymes 2C9, 2C19, and 3A4 with Kvalues (8.5 to 17 mcM) in the range of observed FEV plasma concentrations. In in vitro studies, FEV did not inhibit CYP2E1 and inhibited CYP2D6 and CYP1A2 (K values 82 to 160 mcM). only at concentrations well above those achieved clinically. Coadministration of EFV with drugs primarily metabolized by 2C9, 2C19, and 3A isozymes may result in altered plasma concentrations of the coadministered drug. Drugs which induce CYP3A activity would be expected to Drug interaction studies were performed with FFV and other drugs likely to be coadministered or drugs commonly used as probes for inetic interaction. The effects of coadministration of EFV on the  $C_{max}$ , AUC, and  $C_{min}$  are summarized in Table 8 (effect of EFV on other

drugs) and Table 9 (effect of other drugs on EFV). For information regarding clinical recommendations see Drug Interactions (7.5).

Table 8. Effect of Efavirenz on Coadministered Drug Plasma  $\mathbf{C}_{\scriptscriptstyle{max}},$  AUC, and  $\mathbf{C}_{\scriptscriptstyle{min}}$ 

subjects with hepatic impairment compared with unimpaired subjects.

01	1	ı	Number of		tered Drug (mean	
Coadministered Drug	Dose	Efavirenz Dose	Subjects	C <sub>max</sub> (90% CI)	AUC (90% CI)	C <sub>min</sub> (90% CI)
Boceprevir	800 mg tid x 6 days	600 mg qd x 16 days	NA	↓ 8% (↓ 22 to ↑ 8%)	↓ 19% (11 to 25%)	↓ 44% (26 to 58%)
Simeprevir	150 mg qd x 14 days	600 mg qd x 14 days	23	↓ 51% (↓ 46 to ↓ 56%)	↓ 71% (↓ 67 to ↓ 74%)	↓ 91% (↓ 88 to ↓ 92%)
Ledipasvir/ Sofosbuvir <sup>d</sup>	90/400 mg qd x 14	600 mg qd x 14 days	15	↓ 34	↓ 34	↓ 34
Ledipasvir Sofosbuvir	days			(↓ 25 to ↓ 41) ↔	(↓ 25 to ↓ 41) ↔	(↓ 24 to ↓ 43) NA
GS-331007 <sup>e</sup>				$\leftrightarrow$	$\leftrightarrow$	$\leftrightarrow$
Sofosbuvir <sup>f</sup>	400 mg qd single dose	600 mg qd x 14 days	16	↓ 19 (↓ 40 to ↑10)	$\leftrightarrow$	NA
GS-331007 <sup>e</sup>				↓ 23 (↓ 16 to ↓ 30)	↓ 16 (↓ 24 to ↓ 8)	NA
Sofosbuvir/ Velpatasvir <sup>g</sup>	400/100 mg qd x 14 days	600 mg qd x 14 days	14	(1 10 10 1 30)	(1 24 10 1 0)	
Sofosbuvir	uays			↑ 38 (↑14 to ↑67)	$\leftrightarrow$	NA
GS-331007 <sup>e</sup>				↓ 14 (↓ 20 to ↓ 7)	$\leftrightarrow$	$\leftrightarrow$
Velpatasvir				↓ 47 (↓ 57 to ↓ 36)	↓ 53 (↓ 61 to ↓ 43)	↓ 57 (↓ 64 to ↓ 48)
Azithromycin	600 mg single dose	400 mg qd x 7 days	14	↑ 22% (4 to 42%)	↔	NA NA
Clarithromycin	500 mg q12h x 7	400 mg qd x 7 days	11	↓ 26% (15 to 35%)	↓ 39%	↓ 53% (42 to 63%)
14-OH metabolite	days			↑ 49%	(30 to 46%) ↑ 34%	(42 to 63%) ↑ 26%
Fluconazole	200 mg x 7 days	400 mg qd x 7 days	10	(32 to 69%) ↔	(18 to 53%) ↔	(9 to 45%) ↔
Itraconazole	200 mg q12h x 28 days	600 mg qd x 14 days	18	↓ 37% (20 to 51%)	↓ 39% (21 to 53%)	↓ 44% (27 to 58%)
Hydroxy-itraconazole	-			↓ 35% (12 to 52%)	↓ 37% (14 to 55%)	↓ 43% (18 to 60%)
Posaconazole	400 mg (oral suspension) bid x 10 and 20 days	400 mg qd x 10 and 20 days	11	↓ 45% (34 to 53%)	↓ 50% (40 to 57%)	NA NA
Rifabutin	300 mg qd x 14 days	600 mg qd x 14 days	9	↓ 32% (15 to 46%)	↓ 38% (28 to 47%)	↓ 45% (31 to 56%)
Voriconazole	400 mg po q12h x 1 day, then 200 mg po q12h x 8 days	400 mg qd x 9 days	NA	↓ 61% <sup>a</sup>	↓ 77% <sup>a</sup>	NA NA
	300 mg po q12h days 2 to 7	300 mg qd x 7 days	NA	↓ 36% <sup>b</sup> (21 to 49%)	↓ 55% b (45 to 62%)	NA
	400 mg po q12h days 2 to 7	300 mg qd x 7 days	NA	↑ 23% b	↓ 7% b (↓ 23 to ↑ 13%)	NA
Artemether/ lumefantrine	Artemether 20 mg/ lumefantrine 120	600 mg qd x 26 days	12			
Artemether dihydroartemisinin	mg tablets (6 4-			↓ 21%	↓ 51%	NA NA
	tablet doses over 3 days)			↓ 38%	↓ 46%	
lumefantrine Atorvastatin	10 mg qd x 4 days	600 mg qd x 15 days	14	↔ ↓ 14%	↓ 21% ↓ 43%	NA ↓ 69%
Total active (including				(1 to 26%) ↓ 15%	(34 to 50%)	(49 to 81%) ↓ 48%
metabolites) Pravastatin	40 mg qd x 4 days	600 mg qd x 15 days	13	(2 to 26%) 32%	(21 to 41%)	(23 to 64%)
Simvastatin	40 mg qd x 4 days	600 mg qd x 15 days	14	(↓ 59 to ↑ 12%) ↓ 72%	(26 to 57%)	(0 to 35%)
	40 mg qu x 4 days	000 mg qu x 15 days	14	(63 to 79%)	(62 to 73%)	(20 to 62%)
Total active (including metabolites)				↓ 68% (55 to 78%)	↓ 60% (52 to 68%)	NA <sup>C</sup>
Carbamazepine	200 mg qd x 3 days, 200 mg bid x 3 days,	600 mg qd x 14 days	12	↓ 20% (15 to 24%)	↓ 27% (20 to 33%)	↓ 35% (24 to 44%)
Epoxide metabolite	then 400 mg qd x 29 days			$\leftrightarrow$	$\leftrightarrow$	↓ 13% (↓ 30 to ↑ 7%)
Cetirizine	10 mg single dose	600 mg qd x 10 days	11	↓ 24% (18 to 30%)	$\leftrightarrow$	NA
Diltiazem	240 mg x 21 days	600 mg qd x 14 days	13	↓ 60% (50 to 68%)	↓ 69% (55 to 79%)	↓ 63% (44 to 75%)
Desacetyl diltiazem				↓ 64% (57 to 69%)	↓ 75% (59 to 84%)	↓ 62% (44 to 75%)
N-monodes-methyl diltiazem				↓ 28% (7 to 44%)	↓ 37% (17 to 52%)	↓ 37% (17 to 52%)
Ethinyl estradiol/ Norgestimate	0.035 mg/0.25 mg x 14 days	600 mg qd x 14 days				
Ethinyl estradiol	1		21	$\leftrightarrow$	$\leftrightarrow$	$\leftrightarrow$
Norelgestromine	]		21	↓ 46% (39 to 52%)	↓ 64% (62 to 67%)	↓ 82% (79 to 85%)
Levonorgestrel	1		6	↓ 80% (77 to 83%)	↓ 83% (79 to 87%)	↓ 86% (80 to 90%)
Lorazepam	2 mg single dose	600 mg qd x 10 days	12	↑ 16% (2 to 32%)	↔	NA NA
Methadone	Stable maintenance 35 to 100 mg daily	600 mg qd x 14 to 21 days	11	↓ 45% (25 to 59%)	↓ 52% (33 to 66%)	NA
Bupropion	150 mg single dose	600 mg qd x 14 days	13	↓ 34%	↓ 55%	NA
Hydroxy-bupropion	(sustained-release)			(21 to 47%) ↑ 50%	(48 to 62%) ↔	NA
Paroxetine	20 mg gd x 14 days	600 mg ad v 14 days	16	(20 to 80%)	$\leftrightarrow$	↔

↑ Indicates increase ↓ Indicates decrease ↔ Indicates no change or a mean increase or decrease of < 10%. Relative to steady-state administration of voriconazole (400 mg for 1 day, then 200 mg po q12h for 2 days). Not available because of insufficient data. Study conducted with ATRIPLA®coadministered with HARVONI®

600 mg qd x 14 da

0 mg qd x 14 days | 600 mg qd x 14 days

mg qd x 14 days

\* The predominant circulating nucleoside metabolite of sofosbuvir.

Study conducted with ATRIPLA coadministered with SOVALDI® (sofosbuvir). Study conducted with ATRIPLA coadministered with EPCLUSA® NA = not available.

Table 9. Effect of Coadministered Drug on Efavirenz Plasma  $C_{\mbox{\tiny max}}, AUC, and C_{\mbox{\tiny min}}$ 

Coadministered			l of	•	4110	_	doses up to 4,000 mg per kg per day, pro	oudoing pidoina
Drug	Dose	Efavirenz Dose	Subjects	C <sub>max</sub> (90% CI)	AUC (90% CI)	C <sub>min</sub> (90% CI)	effect on the survival, growth, and develop Tenofovir Disoproxil Fumarate: Long-te	
Boceprevir	800 mg tid x 6 days	600 mg qd x 16 days	NA	↑ 11% (2 to 20%)	↑ 20% (15 to 26%)	NA	16 times (mice) and 5 times (rats) those adenomas were increased at exposures	e observed in hu 16 times that in
Simeprevir	150 mg qd x 14 days	600 mg qd x 14 days	23	$\leftrightarrow$	↓ 10% (5 to 15%)	↓ 13% (7 to 19%)	times that observed in humans at the ther TDF was mutagenic in the <i>in vitro</i> mouse micronucleus assay, TDF was negative wh	lymphoma assa
Azithromycin	600 mg single dose	400 mg qd x 7 days	14	$\leftrightarrow$	$\leftrightarrow$	$\leftrightarrow$	There were no effects on fertility, matin	
Clarithromycin	500 mg q12h x 7 days	400 mg qd x 7 days	12	↑ 11% (3 to 19%)	$\leftrightarrow$	$\leftrightarrow$	equivalent to 10 times the human dose ba mating through day seven of gestation. Th	
Fluconazole	200 mg x 7 days	400 mg qd x 7 days	10	↔	↑ 16%	↑ 22%	13.2 Animal Toxicology and/or Pharma Etavirenz: Nonsustained convulsions we	
Tuconazoic		400 mg qu x 7 days	10	.,	(6 to 26%)	(5 to 41%)	than those in humans given the recomme  Tenofovir Disoproxil Fumarate: Tenofov	nded dose [see l
Itraconazole	200 mg q12h x 14 days	600 mg qd x 28 days	16	$\leftrightarrow$	$\leftrightarrow$	$\leftrightarrow$	greater than or equal to 6-fold those ob- Osteomalacia observed in monkeys app	served in humar
Rifabutin	300 mg qd x 14 days	600 mg qd x 14 days	11	$\leftrightarrow$	$\leftrightarrow$	↓ 12% (↓ 24 to ↑1%)	toxicity manifested as reduced bone mine Evidence of renal toxicity was noted in 4 a	eral density. The r nimal species. Ir
Rifampin	600 mg x 7 days	600 mg qd x 7 days	12	↓ 20% (11 to 28%)	↓ 26% (15 to 36%)	↓ 32% (15 to 46%)	and decreases in serum phosphate were on to 20 times higher than those observed in not known.	
Voriconazole	400 mg po q12h x 1 day, then 200 mg po	400 mg qd x 9 days	NA	↑ 38% <sup>a</sup>	↑ 44% <sup>a</sup>	NA	14 CLINICAL STUDIES	
	q12h x 8 days						14.1 Clinical Efficacy in Patients with H	
	300 mg po q12h days 2 to 7	300 mg qd x 7 days	NA	↓ 14% <sup>b</sup> (7 to 21%)	<b>b</b> ↔	NA	Treatment-Naïve Adult Patients: The eff antiretroviral treatment history was estab • Trial 903 which evaluated the efficar	lished in trials of cy of a three-drug
	400 mg po q12h days 2 to 7	300 mg qd x 7 days	NA	↔b	↑ 17% <sup>b</sup> (6 to 29%)	NA	<ul> <li>ENCORE1, which evaluated the con Trial 903: Data through 144 weeks are re mg + TDF 300 mg vs. EFV 600 mg + 3TC 3</li> </ul>	ported for Trial 9
Artemether/ Lumefantrine	Artemether 20 mg/ lumefantrine 120 mg tablets (6 4-tablet doses over 3 days)	600 mg qd x 26 days	12	↔	↓ 17%	NA	(range 18 to 64); 74% were male, 64% v 956) and median baseline plasma HIV-1 CD4+ cell count. Forty-three percent of su 11 provides treatment outcomes through Table 11. Outcomes of Randomized Tre	vere Caucasian, RNA was 77,600 Ibjects had baseli 48 and 144 weel
Atorvastatin	10 mg qd x 4 days	600 mg qd x 15 days	14	$\leftrightarrow$	$\leftrightarrow$	$\leftrightarrow$	Outcomes	atiliont at wook
Pravastatin	40 mg qd x 4 days	600 mg qd x 15 days	11	↔	$\leftrightarrow$	$\leftrightarrow$	Guicomes	EFV + 3TC - (N = 29
Simvastatin	40 mg qd x 4 days	600 mg qd x 15 days	14	↓ 12%	$\leftrightarrow$	↓ 12% (↓ 05 to ±20()	Responder <sup>a</sup>	79%
				(↓ 28 to ↑ 8%)		(↓ 25 to ↑3%)	Virologic failure <sup>b</sup>	6%
Aluminum hydroxide 400	30 mL single dose	400 mg single dose	17	$\leftrightarrow$	$\leftrightarrow$	NA	Rebound	5%
mg, magnesium hydroxide 400 mg,							Never suppressed	0%
plus simethicone 40 mg							Added an antiretroviral agent	1%
Carbamazepine	200 mg qd x 3	600 mg qd x 35 days	14	↓ 21%	↓ 36%	↓ 47%	Death	< 1%
	days, 200 mg bid x 3 days, then 400 mg qd x 15 days			(15 to 26%)	(32 to 40%)	(41 to 53%)	Discontinued due to adverse event  Discontinued for other reasons	6% 8%
Cetirizine	10 mg single dose	600 mg qd x 10 days	11	↔	↔	$\leftrightarrow$	Discontinued for other reasons	0 70
Diltiazem	240 mg x 14 days	600 mg qd x 28 days	12	↑ 16% (6 to 26%)	↑ 11% (5 to 18%)	↑ 13% (1 to 26%)	<sup>a</sup> Subjects achieved and maintained confir <sup>b</sup> Includes confirmed viral rebound and fai <sup>c</sup> Includes lost to follow-up, subject's with	ilure to achieve c
Famotidine	40 mg single dose	400 mg single dose	17	(0 (0 20 /0)	(3 t0 10 /0)	(1 to 20 %)	Achievement of plasma HIV-1 RNA con	centrations of <
Paroxetine	20 mg qd x 14 days	600 mg qd x 14 days	12	↔	↔	+ H	population stratified at baseline on the ba	
Sertraline	50 mg qd x 14 days	600 mg qd x 14 days	13	↑ 11%	↔	↔	Through 144 weeks of therapy, 62% and < 50 copies/mL. The mean increase from	
	,			(6 to 16%)			Through 144 weeks, 11 subjects in the TD	F group and 9 su

Efavirez (mean % change)

 $Indicates\ increase \downarrow Indicates\ decrease \leftrightarrow Indicates\ no\ change\ or\ a\ mean\ increase\ or\ decrease\ of\ <10\%.$ 90% CI not available Relative to steady-state administration of efavirenz (600 mg once daily for 9 days).

NA = not available. Lamivudine: Effect of 3TC on the Pharmacokinetics of Other Agents: Based on in vitro study results, 3TC at therapeutic drug exposures is not expected to affect the pharmacokinetics of drugs that are substrates of the following transporters: organic anion transporter polypeptide 1B1/3 (OATP1B1/3), breast cancer resistance protein (BCRP), P-glycoprotein (P-gp), multidrug and toxin extrusion protein 1 (MATE1), MATE2-K, organic cation transporter 1 (OCT1), OCT2, or OCT3. Effect of Other Agents on the Pharmacokinetics of 3TC: 3TC is a substrate of MATE1, MATE2- K, and OCT2 in vitro. Trimethoprim (an inhibitor of

adjustment of 3TC is needed. 3TC is a substrate of P-gp and BCRP; however, considering its absolute bioavailability (87%), it is unlikely that these transporters play a significant ole in the absorption of 3TC. Therefore, coadministration of drugs that are inhibitors of these efflux transporters is unlikely to affect the disp

Interferon Alfa: There was no significant pharmacokinetic interaction between 3TC and interferon alfa in a trial of 19 healthy male subjects. Ribavirin: In vitro data indicate ribavirin reduces phosphorylation of 3TC, stayudine, and zidovudine. However, no pharmacokinetic (e.g., plasma concentrations or intracellular triphosphorylated active metabolite concentrations) or pharmacodynamic (e.g., loss of HIV-1/HCV virologic suppression) interaction was observed when ribavirin and 3TC (n = 18), stavudine (n = 10), or zidovudine (n = 6) were coadministered as part of a multi-drug regimen to HIV-1/HCV co-infected subjects. Sorbitol (Ex<u>cipient):</u> 3TC and sorbitol solutions were coadministered to 16 healthy adult subjects in an open-label, randomized-sequence, 4-perio

crossover trial. Each subject received a single 300-mg dose of 3TC oral solution alone or coadministered with a single dose of 3.2 grams, 10.2 rrams or 13.4 grams of sorbitol in solution. Coadministration of 3TC with sorbitol resulted in dose-dependent decreases of 20%, 39%, and 44% in the AUC  $_{(00024)}$ , 14%, 32%, and 36% in the AUC  $_{(\omega)}$ , and 28%, 52%, and 55% in the C $_{max}$  of lamivudine, respectively. Frimethoprim/Sulfamethoxazole: 3TC and TMP/SMX were coadministered to 14 HIV-1-positive subjects in a single-center, open-label, t with a single 300-mg dose of 3TC and TMP 160 mg/SMX 800 mg once a day for 5 days with concomitant administration of 3TC 300 mg with the fifth dose in a crossover design. Coadministration of TMP/SMX with 3TC resulted in an increase of  $43\% \pm 23\%$  (mean  $\pm$  SD) in 3TC AUC. a decrease of  $29\% \pm 13\%$  in 3TC oral clearance, and a decrease of  $30\% \pm 36\%$  in 3TC renal clearance. The pharmacokinetic properties of TMP and SMX were not altered by coadministration with 3TC. There is no information regarding the effect on 3TC pharmacokinetics of higher doses of TMP/SMX such as those used in treating PCP.

CYP3A4, CYP2D6, CYP2C9, or CYP2E1. However, a small (6%) but statistically significant reduction in metabolism of CYP1A substrate was observed. Based on the results of in vitro experiments and the known elimination pathway of tenofovir, the potential for CYP-mediated interactions Table 10 summarizes pharmacokinetic effects of coadministered drug on tenofovir pharmacokinetics. No clinically significant drug interactions have been observed between tenofovir and ribavirin.

Table 10. Drug Interactions: Changes in Pharmacokinetic Parameters for Tenofovir<sup>a</sup>in the Presence of the Coadmini

Tenofovir Disoproxil Fumarate: At concentrations substantially higher (~300-fold) than those observed in vivo tenofovir did not inhibit in vitro

Table 10. Drug Iliteraction	ns: Gnanges in Pharmacokinetic Parame	515131			
Coadministered Drug	Dose of Coadministered Drug (mg)		% Change of T	enofovir Pharmacokinetic (90% CI)	Parameters <sup>b</sup>
			C <sub>max</sub>	AUC	C <sub>min</sub>
Ledipasvir/ Sofosbuvir <sup>e,f</sup>	90/400 once daily x 10 days	24	↑ 47 (↑ 37 to ↑ 58)	↑ 35 (↑ 29 to ↑ 42)	↑ 47 (↑ 38 to ↑ 57)
Ledipasvir/ Sofosbuvir <sup>e,g</sup>		23	↑ 64 (↑ 54 to ↑ 74)	↑ 50 (↑ 42 to ↑ 59)	↑ 59 (↑ 49 to ↑ 70)
Ledipasvir/ Sofosbuvir <sup>c</sup>	90/400 once daily x 14 days	15	↑ 79 (↑ 56 to ↑104)	↑ 98 (↑ 77 to ↑123)	↑ 163 (↑ 132 to ↑ 197)
Sofosbuvir/ Velpatasvir <sup>h</sup>	400/100 once daily	24	↑ 44 (↑ 33 to ↑55)	↑ 40 (↑ 34 to↑46)	↑ 84 (↑ 76 to ↑ 92)
Sofosbuvir/ Velpatasvir <sup>i</sup>	400/100 once daily	30	↑ 46 (↑ 39 to ↑54)	↑ 40 (↑ 34 to ↑45)	↑70 (↑61 to ↑ 79)
Sofosbuvir/ Velpatasvir/ Voxilaprevir <sup>i</sup>	400/100/100 + Voxilaprevir* 100 once daily	29	↑ 48 (↑ 36 to ↑61)	↑ 39 (↑ 32 to ↑46)	↑47 (↑38 to ↑ 56)
Sofosbuvir <sup>d</sup>	400 single dose	16	↑ 25 (↑ 8 to ↑ 45)	$\leftrightarrow$	$\leftrightarrow$
Tacrolimus	0.05 mg/kg twice daily x 7 days	21	↑ 13 (↑ 1 to ↑ 27)	$\leftrightarrow$	$\leftrightarrow$

Subjects received tenofovir disoproxil furnarate 300 mg once daily  $Increase = \uparrow$ ;  $Decrease = \downarrow$ ;  $No Effect = \leftrightarrow$ ; NC = Not CalculatedStudy conducted with efavirenz/emtricitabine/tenofovir disoproxil fumarate coadministered with ledipasyir/sofosbuyi

of mammalian DNA polymerases  $\alpha$ ,  $\beta$ , and mitochondrial DNA polymerase  $\gamma$ .

Study conducted with efavirenz/emtricitabine/tenofovir disoproxil fumarate coadministered with sofosbuving Data generated from simultaneous dosing with ledipasvir/sofosbuvir. Staggered administration (12 hours apart) provide similar results. Comparison based on exposures when administered as atazanavir/ritonavir + emtricitabine/tenofovir DF Comparison based on exposures when administered as darunavir/ritonavir + emtricitabine/tenofovir DF Study conducted with COMPLERA coadministered with EPCLUSA; coadministration with EPCLUSA also results in comparable increases in

tenofovir exposures when TDF is administered as ATRIPLA, STRIBILD® (elvitegravir/cobicistat/FTC/TDF), TRUVADA + atazanavir/ritonavir, or Administered as raltegravir + FTC/TDF. res when administered as darunavir + ritonavir + FTC/TDF. Study conducted with additional voxilaprevir 100 mg to achieve voxilaprevir exposures expected in HCV-infected patients

Mechanism of Action: Efavirenz: EFV is an NNRTI of HIV-1. EFV activity is mediated predominantly by noncompetitive inhibition of HIV-1 reverse  $transcript as e \, (RT). \, HIV-2 \, RT \, and \, human \, cellular \, DNA \, polymerases \, \alpha, \, \beta, \, \gamma, \, and \, \delta \, are \, not \, inhibited \, by \, EFV.$ Lamivudine: 3TC is a synthetic nucleoside analogue with activity against HIV-1 and HBV. Intracellularly, 3TC is phosphorylated to its active 5'-triphosphate metabolite, lamivudine triphosphate (3TC-TP). The principal mode of action of 3TC-TP is inhibition of HIV-1 reverse transcriptase (RT) via DNA chain termination after incorporation of the nucleotide analogue. Tenofovir Disoproxil Fumarate: TDF is an acyclic nucleoside phosphonate diester analog of adenosine monophosphate. TDF requires initial diester hydrolysis for conversion to tenofovir and subsequent phosphorylations by cellular enzymes to form tenofovir diphosphate (TDF-DP), an obligate chain terminator. Tenofovir diphosphate inhibits the activity of HIV-1 reverse transcriptase (RT) and HBV RT by competing with the natural substrate deoxyadenosine 5'-triphosphate and, after incorporation into DNA, by DNA chain termination. Tenofovir diphosphate is a weak inhibitor

Antiviral Activity: Efavirenz: The concentration of EFV inhibiting replication of wild-type laboratory adapted strains and clinical isolates in cell culture by 90 to 95% (EC $_{\infty}$   $_{\infty}$   $_{\infty}$ ) ranged from 1.7 to 25 nM in lymphoblastoid cell lines, peripheral blood mononuclear cells (PBMCs), and onocyte cultures. EFV demonstrated antiviral activity against clade B and most non-clade B isolates (subtypes A, AE, AG, C, D, F, G, J, N), but had reduced antiviral activity against group O viruses. Lamivudine: The antiviral activity of 3TC against HIV-1 was assessed in a number of cell lines (including monocytes and fresh human peripheral blood lymphocytes (PBMCs) using standard susceptibility assays. EC $_{so}$  values were in the range of 3 to 15,000 n/M. (1 mcM = 0.23 mcg/mL). The median EC $_{so}$  values of 3TC were 60 nM (range: 20 to 70 nM), 35 nM (range: 30 to 40 nM), 30 nM (range: 20 to 90 nM), 20 nM (range: 3 to 40 nM), 30 nM (range: 20 to 90 nM). nM (range: 1 to 60 nM), 30 nM (range: 20 to 70 nM), 30 nM (range: 3 to 70 nM), and 30 nM (range: 20 to 90 nM) against HIV-1 clades A-G and group O viruses (n = 3 except n = 2 for clade B), respectively. The EC<sub>so</sub>values against HIV-2 isolates (n = 4) ranged from 3 to 120 nM in PBMCs. 3TC was not Intagonistic to all tested anti-HIV agents. Ribavirin (50 mcM) used in the treatment of chronic HCV infection decreased the anti-HIV-1 activity of 3TC by 3.5-fold in MT-4 cells.

Tenofovir Disoproxil Fumarate: The antiviral activity of tenofovir against laboratory and clinical isolates of HIV-1 was assessed in lymphoblasto cell lines, primary monocyte/macrophage cells and peripheral blood lymphocytes. The EC on 50% effective concentration) values for tenofovir were in the range of 0.04 mcM to 8.5 mcM. Tenofovir displayed antiviral activity in cell culture against HIV-1 clades A, B, C, D, E, F, G, and O (EC<sub>so</sub> values ranged from 0.5 mcM to 2.2 mcM) and strain-specific activity against HIV-2 (EC<sub>so</sub> values ranged from 1.6 mcM to 5.5 mcM). Please see the full rescribing information for VIREAD $^{\circ}$  for information regarding the inhibitory activity of TDF against HBV. Resistance: Efavirenz: In cell culture, HIV-1 isolates with reduced susceptibility to EFV (> 380-fold increase in EC to 2014 and 2014 energed rapidly in the presence of drug. Genotypic characterization of these viruses identified single amino acid substitutions L100l or V179D, double substitutions

L100I/V108I, and triple substitutions L100I/V179D/Y181C in reverse transcriptase. Clinical isolates with reduced susceptibility in cell culture to EFV have been obtained. One or more RT substitutions at amino acid positions A98, L100, K101, K103, V106, V108, Y188, G190, P225, F227 and M230 were observed in patients failing treatment with EFV in combination with ndinavir, or with 3TC plus zidovudine. The K103N substitution was the most frequently observed. Lamivudine: 3TC-resistant variants of HIV-1 have been selected in cell culture. Genotypic analysis showed that resistance was predominantly due to a methionine to valine or isoleucine (M184V/I) substitution in reverse transcriptase.

Tenofovir Disoproxil Fumarate: HIV-1 isolates with reduced susceptibility to tenofovir have been selected in cell culture. These viruses expressed a K65R substitution in RT and showed a 2- to 4-fold reduction in suscentibility to tenofovir. In addition, a K70F substitution in HIV-1 RT has been selected by tenofovir and results in low-level reduced susceptibility to tenofovir. K65R substitutions developed in some subjects failing a tenofovir disoproxil fumarate regimen. Cross-Resistance: Ffavirenz: Cross-resistance among NNRTIs has been observed. Clinical isolates previously characterized as FFV-resistant were also phenotypically resistant in cell culture to delavirdine and nevirapine compared to baseline. Delavirdine- and/or nevirapine-res viral isolates with NNRTI resistance-associated substitutions (A98G. L100I, K101E/P, K103N/S, V106A, Y181X, Y188X, G190X, P225H, F227L, or M230L) showed reduced susceptibility to EFV in cell culture. Greater than 90% of NRTI-resistant clinical isolates tested in cell culture retained

susceptibility to EFV. Lamivudine: Cross-resistance among NRTIs has been observed. 3TC-resistant HIV-1 isolates were cross-resistant in cell culture to didanosine ddl). Cross-resistance is also expected with abacavir and emtricitabine as these select M184V substitution Tenofovir Disoproxil Fumarate: Cross-resistance among certain HIV-1 NRTIs has been observed. The K65R and K70E substitutions selected by tenofovir are also selected in some HIV-1-infected subjects treated with abacavir or didanosine. HIV-1 isolates with the K65R substitution also howed reduced susceptibility to FTC and 3TC. HIV-1 isolates from subjects (N = 20) whose HIV-1 expressed a mean of 3 zidovudine-associated RT amino acid substitutions (M41L, D67N, K70R, L210W, T215Y/F, or K219Q/E/N) showed a 3.1-fold decrease in the susceptibility to tenofovir. Subjects whose virus expressed an L74V substitution without zidovudine resistance-associated substitutions (N = 8) had reduced response to VIREAD. Limited data are available for patients whose virus expressed at Y115F substitution (N = 3), Q151M substitution (N = 2), or T69 insertion (N = 4), all of whom had a reduced response

13 NONCLINICAL TOXICOLOGY 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Elavirenz: Long-term carcinogenicity studies in mice and rats were carried out with efavirenz. Mice were dosed with 0, 25, 75, 150, or 300 mg/kg/day for 2 years. Incidences of hepatocellular adenomas and carcinomas and pulmonary alveolar/bronchiolar adenomas were increased bove background in females. No increases in tumor incidence above background were seen in males. There was no NOAEL in female established for this study because tumor findings occurred at all doses. AUC at the NOAEL (150 mg/kg) in the males was approximately 0.9 times that in ended clinical dose. In the rat study, no increases in tumor incidence were observed at doses up to 100 mg/kg/day, for which AUCs were 0.1 (males) or 0.2 (females) times those in humans at the recommended clinical dose. EFV tested negative in a battery of in vitro and in vivo genotoxicity assays. These included bacterial mutation assays in S. typhimurium and E. coli, mammalian mutation assays in Chinese hamster ovary cells, chromosome aberration assays in human peripheral blood lymphocytes or Chinese hamster ovary cells, and an *in vivo* mouse bone marrow micronucleus assay.

EFV did not impair mating or fertility of male or female rats, and did not affect sperm of treated male rats. The reproductive performance of offspring oorn to female rats given EFV was not affected. The AUCs at the NOAEL values in male (200 mg/kg) and female (100 mg/kg) rats were approximately < 0.15 times that in humans at the recommended clinical dose. Lamivudine: Long-term carcinogenicity studies with 3TC in mice and rats showed no evidence of carcinogenic potential at exposures up to 10

assay, in an in vitro cell transformation assay, in a rat micronucleus test, in a rat bone marrow cytogenetic assay, and in assay for unscheduled DNA synthesis in rat liver. 3TC showed no evidence of in vivo genotoxic activity in the rat at oral doses of up to 2,000 mg per kg, producing plasma levels of 35 to 45 times those in humans at the recommended dose for HIV-1 infection. In a study of reproductive performance, 3TC admir doses up to 4.000 mg per kg per day, producing plasma levels 47 to 70 times those in humans, revealed no evidence of impaired fertility and no

 $\hbox{\it effect on the survival, growth, and development to weaning of the offspring.}$ Tenofovir Disoproxil Fumarate: Long-term oral carcinogenicity studies of TDF in mice and rats were carried out at exposures up to approxi 16 times (mice) and 5 times (rats) those observed in humans at the therapeutic dose for HIV-1 infection. At the high dose in female mice, liver mas were increased at exposures 16 times that in humans. In rats, the study was negative for carcinogenic findings at exposures up to 5 times that observed in humans at the therapeutic dose. TDF was mutagenic in the in vitro mouse lymphoma assay and negative in an in vitro bacterial mutagenicity test (Ames test). In an in vivo mouse

nucleus assay, TDF was negative when administered to male i There were no effects on fertility, mating performance or early embryonic development when TDF was administered to male rats at a dose equivalent to 10 times the human dose based on body surface area comparisons for 28 days prior to mating and to female rats for 15 days prior to mating through day seven of gestation. There was, however, an alteration of the estrous cycle in female rats.

13.2 Animal Toxicology and/or Pharmacology Elavirenz: Nonsustained convulsions were observed in 6 of 20 monkeys receiving EFV at doses yielding plasma AUC values 4- to 13-fold greater than those in humans given the recommended dose [see Warnings and Precautions (5.11)]. Tenofovir Disoproxil Fumarate: Tenofovir and TDF administered in toxicology studies to rats, dogs, and monkeys at exposures (based on AUCs) greater than or equal to 6-fold those observed in humans caused bone toxicity. In monkeys the bone toxicity was diagnosed as osteomalacia. Osteomalacia observed in monkeys appeared to be reversible upon dose reduction or discontinuation of tenofovir. In rats and dogs, the bone

oxicity manifested as reduced bone mineral density. The mechanism(s) underlying bone toxicity is unknown Evidence of renal toxicity was noted in 4 animal species. Increases in serum creatinine, BUN, glycosuria, proteinuria, phosphaturia, and/or calciuria and decreases in serum phosphate were observed to varying degrees in these animals. These toxicities were noted at exposures (based on AUCs) 2 to 20 times higher than those observed in humans. The relationship of the renal abnormalities, particularly the phosphaturia, to the bone toxicity is 14 CLINICAL STUDIES

Treatment-Naïve Adult Patients: The efficacy of EFV 400 mg, 3TC 300 mg, and TDF 300 mg in the treatment of HIV-1 infection in adults with no antiretroviral treatment history was established in trials of:
Trial 903 which evaluated the efficacy of a three-drug regimen including EFV 600 mg, 3TC 300 mg and TDF 300 mg ENCORE1, which evaluated the comparability of 400 mg of EFV in a triple drug regimen to a 600 mg dose of EFV in a triple drug regimen. Trial 903: Data through 144 weeks are reported for Trial 903, a double-blind, active-controlled multicenter trial comparing EFV 600 mg + 3TC 300 mg + TDF 300 mg vs. EFV 600 mg + 3TC 300 mg + stavudine (d4T) 40 mg in 600 antiretroviral-naïve subjects. Subjects had a mean age of 36 years (range 18 to 64); 74% were male, 64% were Caucasian, and 20% were Black. The mean baseline CD4+ cell count was 279 cells/mm² (range 3 to 956) and median baseline plasma HIV-1 RNA was 77,600 copies/mL (range 417 to 5,130,000). Subjects were stratified by baseline HIV-1 RNA and

Outcomes	At We	eek 48	At W	eek 144
	EFV + 3TC + TDF (N = 299)	EFV + 3TC + d4T (N = 301)	EFV + 3TC + TDF (N = 299)	EFV + 3TC + d4T (N = 301)
Responder <sup>a</sup>	79%	82%	68%	62%
Virologic failure <sup>b</sup>	6%	4%	10%	8%
Rebound	5%	3%	8%	7%
Never suppressed	0%	1%	0%	0%
Added an antiretroviral agent	1%	1%	2%	1%
Death	< 1%	1%	< 1%	2%
Discontinued due to adverse event	6%	6%	8%	13%
Discontinued for other reasons <sup>c</sup>	8%	7%	14%	15%

<sup>a</sup> Subjects achieved and maintained confirmed HIV-1 RNA < 400 copies/mL through Week 48 and 144. Includes confirmed viral rebound and failure to achieve confirmed < 400 copies/mL through Week 48 and 144. Includes lost to follow-up, subject's withdrawal, noncompliance, protocol violation and other reasons.

RNA level, either < 100,000 copies/mL or ≥ 100,000 copies/ml

Achievement of plasma HIV-1 RNA concentrations of < 400 copies/mL at Week 144 was similar between the two treatment groups for the population stratified at baseline on the basis of HIV-1 RNA concentration (> or ≤ 100.000 copies/mL) and CD4+ cell count (< or ≥ 200 cells/mm³). hrough 144 weeks of therapy, 62% and 58% of subjects in the TDF and d4T arms, respectively, achieved and maintained cont < 50 copies/mL. The mean increase from baseline in CD4+ cell count was 263 cells/mm³ for the TDF arm and 283 cells/mm³ for the d4T arm. Through 144 weeks, 11 subjects in the TDF group and 9 subjects in the d4T group experienced a new CDC Class C event. The ENCORE1 trial was a randomized, multinational clinical study comparing EFV 400 mg vs. EFV 600 mg in 630 antiretroviral-naïve adult subjects. Subjects were randomized 1:1 to receive EFV 400 mg in combination with TDF 300 mg plus FTC 200 mg all given once daily or EFV 600 mg in

Subjects had a mean age of 36 years (range 18 to 69), 68% were male, 37% were of African heritage, 33% were of Asian ethnicity, 17% were Hispanic and 13% were Caucasian The mean baseline CD4+ cell count was 273 cells/mm³ (range 38 to 679). Median baseline viral load was 56,469 copies/mL (range 162 to 10,000,000). Thirty-four percent of subjects had baseline viral load of  $\geq$  100,000 copies/mL. Treatment outcomes through Week 48 are presented in Table 12.

ombination with TDF 300 mg/FTC 200 mg given once daily. The randomization was stratified by the clinical sites and the screening visit plasma HIV

Table 12. Virologic Outcomes of Randomized Treatment in Trial ENCORE1 in Treatment-Naïve Subjects at Week 48

	At Week 48					
Outcomes (< 50 copies/mL)	EFV 400 mg + FTC + TDF (N = 321)	EFV 600 mg + FTC + TDF (N = 309)				
Responder <sup>a</sup> HIV-1 RNA < 50 copies/mL	86%	84%				
Virologic failure <sup>b</sup> HIV-1 RNA ≥ 50 copies/mL	11%	11%				
Rebound	9%	8%				
Never suppressed	2%	3%				
Death	1%	1%				
Discontinued for other reasons <sup>c</sup>	2%	4%				

\*Includes confirmed viral rebound and failure to achieve confirmed <50 copies/mL through Week 48.

\*Includes discontinued due to Adverse Event, lost to follow-up, subject's withdrawal, noncompliance, protocol violation and other reasons. Achievement of plasma HIV-1 RNA concentrations of less than 50 copies/mL at Week 48 was similar between the two treatment groups for the ulation stratified at baseline on the basis of HIV-1 RNA concentration (< or ≥ 100,000 copies/mL). The mean increase at Week 48 from baseline in CD4 $^{\circ}$ cell count was 183 cells/mm $^{\circ}$  for the EFV 400 mg arm and 158 cells/mm $^{\circ}$  for the EFV 600 mg arm. Through 48 weeks, 11 subjects in the EFV 400 mg group and 5 subjects in the EFV 600 mg group experienced a new CDC Class C event. 16 HOW SUPPLIED/STORAGE AND HANDLING

Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are supplied as fixed-dose combination tablets containing efavirenz USP 400 mg, Lamivudine USP 300 mg and tenofovir disoproxii fumarate 300 mg equivalent to tenofovir disoproxii 245 mg. Efavirenz, lamivudine and tenofovir disoproxii fumarate 300 mg equivalent to tenofovir disoproxii 245 mg. Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are white to off-white, film-coated, oval, biconvex tablets debossed with "L40" on one side and plain on other side. They are supplied as follows: HDPE bottles with Molecular sieve desiccant

NDC 42385-929-30 Bottles of 90 NDC 42385-929-90 Bottles of 180 NDC 42385-929-18 HDPE bottles with Silica gel canister desiccant NDC 42385-929-31 Bottles of 30 Bottles of 90 Bottles of 180 NDC 42385-929-82 Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.]

17 PATIENT COUNSELING INFORMATION  $Advise the \ patient to \ read \ the \ FDA-approved \ patient \ labeling \ (Patient \ Information).$ 

**Drug Interactions:** Efavirenz, lamivudine and tenofovir disoproxil fumarate may interact with many drugs; therefore, advise patients to report to their healthcare provider the use of any other prescription, nonprescription medication, or herbal products, particularly St. John's wort *Isee* Contraindications (4) and Drug Interactions (7)]. Post Treatment Acute Exacerbation of Hepatitis B in Patients with HBV Co-Infection: Inform patients that severe acute exacerbations of hepatitis have been reported in patients who are infected with HBV or coinfected with HBV and HIV-1 and have discontinued 3TC and TDF, components of

efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Test patients with HIV-1 for hepatitis B virus (HBV) before initiating antiretroviral therapy. In patients with chronic hepatitis B, it is important to obtain HIV antibody testing prior to initiating 3TC and TDF, components of efavirenz, [5.1) lamivudine and tenofovir disoproxil fumarate tablets [see Warnings and Precautions Lactic Acidosis and Severe Hepatomegaly: Inform patients that lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported. Treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate should be suspended in any patient who develops clinical symptoms suggestive of lactic acidosis or pronounced hepatotoxicity (including nausea, vomiting, unusual or unexpected stomach discomfort, and weakness) [see Warnings and Precautions (5.2)]. New Onset or Worsening Renal Impairment: Inform patients that renal impairment, including cases of acute renal failure and Fanconi syndrome,

has been reported. Advise patients with impaired renal function (i.e., creatinine clearance less than 50 mL/min) or patients with end-stage renal disease (ESRD) requiring hemodialysis to avoid efavirenz, lamivudine and tenofovir disoproxil fumarate with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple NSAIDs) for patients [see Dosage and Administration (2.3), Warnings and Precautions (5.4)]. Psychiatric Symptoms: Inform patients that serious psychiatric symptoms including severe depression, suicide attempts, aggressive behavior delusions, paranoia, psychosis-like symptoms and catatonia have been reported in patients receiving EFV [see Warnings and Precautions (5.5)]. Advise patients to seek immediate medical evaluation if they experience severe psychiatric adverse experiences. Advise patients to inform their physician of any history of mental illness or substance abuse. Nervous System Symptoms: Inform patients that central nervous system symptoms (NSS) including dizziness, insomnia, impaired concentratio drowsiness, and abnormal dreams are commonly reported during the first weeks of therapy with EFV, a component of efavirenz, lamivudine and

tenofovir disoproxil furnarate tablets [see Warnings and Precautions (5.6)]. Dosing at bedtime may improve the tolerability of these symptoms, which are likely to improve with continued therapy. Alert patients to the potential for additive effects when used concomitantly with alcohol or

psychoactive drugs. Instruct patients that if they experience NSS they should avoid potentially hazardous tasks such as driving or operating

machinery. Inform patients that there is a risk of developing late-onset neurotoxicity, including ataxia and encephalopathy, which may occur months to years after beginning efavirenz therapy [see Warnings and Precautions (5.6)]. Embryo-Fetal Toxicity: Advise female patients that EFV, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets may cause fetal harm when administered during the first trimester to a pregnant woman. Advise females of reproductive potential to use effective contraception as well as a barrier method during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate and for 12 weeks after discontinuation of use. Advise patients to contact their healthcare provider if they plan to become pregnant, become pregnant, or if pregnancy is suspected during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate [see Warnings and Precautions (5.7), Use in Specific

Populations (8.1, 8.3)]. Rash: Inform patients that rash is a common side effect of EFV [see Warnings and Precautions (5.8)]. Rashes usually go away without any change in treatment. However, since rash may be serious, patients should be advised to contact their physician promptly if rash occurs Hepatotoxicity: Inform patients to watch for early warning signs of liver inflammation or failure, such as fatigue, weakness, lack of appetite, nausea and vomiting, as well as later signs such as jaundice, confusion, abdominal swelling, and discolored feces and to consult their healthcare provider promptly if such symptoms occur [see Warnings and Precautions (5.9)].

Pancreatitis: Advise patients or guardians to monitor pediatric patients for signs and symptoms of pancreatitis [see Warnings and Precautions Convulsions: Advise patients that convulsions have been observed in patients receiving EFV, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, generally in patients with known medical history of seizures [see Warnings and Precautions (5.11)]. Lipid Elevations: Advise patients treatment with EFV, a component of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets has resulted in es in the concentration of total cholesterol and triglycerides [see Warnings and Precautions (5.12)]. Bone Loss and Mineralization Effects: Inform patients that decreases in bone mineral density have been observed with the use of 3TC and TDF, components of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, in patients with HIV [see Warnings and Precautions (5.13)].

Immune Reconstitution Syndrome: Advise patients to inform their healthcare provider immediately of any symptoms of infection, as in some

atients with advanced HIV infection, signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is

Fat Redistribution: Inform patients that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy, including virenz, lamivudine and tenofovir disoproxil fumarate, and that the cause and long-term health effects of these conditions are not known at this time [see Warnings and Precautions (5.15)]. Administration Instructions: Inform patients that it is important to take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets once daily on a regular dosing schedule on an empty stomach, preferably at bedtime, and to avoid missing doses as it can result in development of resistance. Advise patients if a dose is missed, take it as soon as possible unless it is almost time for the next dose. Also advise patients that dosing at bedtime may improve the tolerability of nervous system symptoms [see Dosage and Administration (2.2)].

**Pregnancy Registry:** Advise patients that there is an antiretroviral pregnancy registry to monitor fetal outcomes in women exposed to efavirenz, lamivudine and tenofovir disoproxil fumarate [see Use in Specific Populations (8.1)]. Lactation: Instruct women with HIV-1 infection not to breastfeed because HIV-1 can be passed to the baby in breast milk [see Use in Specific Populations (8.2)1. Brands listed are the trademarks of their respective owners and are not trademarks of Laurus Labs Limited.

Laurus Generics Inc 400 Connell Drive Suite 5200 Berkeley Heights, NJ 07922 Manufactured by:

Anakapalli-531011

India

started [see Warnings and Precautions (5.14)].

## **Patient Information** Efavirenz, Lamivudine and Tenofovir Disoproxil Fumarate (ef-a-vir-enz, la-MIV-ue-deen and ten-OF-oh-vir DYE-soe-PROX-il FUE-ma-rate) Tablets

What is the most important information I should know about efavirenz, lamivudine and tenofovir disoproxil fumarate tablets?

 $\textbf{Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets can cause serious side\ effects, including: \\$ Worsening of Hepatitis B virus infection. If you have Human Immunodeficiency Virus type 1 (HIV-1) and Hepatitis B Virus (HBV) infection

your HBV may get worse (flare-up) if you stop taking efavirenz, lamiyudine and tenofovir disporoxil fumarate tablets. A "flare-up" is when our HBV infection suddenly returns in a worse way than before. Your healthcare provider will test you for HBV infection before you start treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. It is not known if efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are safe and effective in people who have both HIVand HBV infection.

Do not run out of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Refill your prescription or talk to your healthcare provider before your efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are all gone. Do not stop efavirenz, lamivudine and tenofovir disoproxil fumarate tablets without first talking to your healthcare provider. If you stop taking efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, your healthcare provider will need to check your health often and do blood tests regularly for several months to check your liver.

For more information about side effects, see "What are the possible side effects of efavirenz, lamivudine and tenofovir disoproxil fumarate What are efavirenz, lamivudine and tenofovir disoproxil fumarate tablets? Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets is a prescription medicine that is used without other antiretroviral medicines to

treat Human Immunodeficiency Virus-1 (HIV-1) in people weighing at least 35 kg. HIV-1 is the virus that causes AIDS (Acquired Immune Deficiency Syndrome) Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets contains the prescription medicines efavirenz, lamivudine and tenofovir favirenz, lamivudine and tenofovir disoproxil fumarate tablets are not for use in children weighing less than 35 kg.

Do not take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets if you: are allergic to efavirenz, lamivudine, tenofovir disoproxil fumarate, or any of the ingredients in efavirenz, lamivudine and tenofovir disoproxi fumarate tablets. See the end of this Patient Information leaflet for a complete list of ingredients in efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

are currently taking elbasvir and grazoprevir Before you take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, tell your healthcare provider about all of your medical

ditions, including if you:

have liver problems, including hepatitis B or C infection have kidney problems, including end-stage renal disease (ESRD) that requires dialysis

have a history of mental health problems have a history of drug or alcohol abuse have a heart problem, including QT prolongation

have bone problems, including a history of bone fractures

are pregnant or plan to become pregnant. Efavirenz, lamivudine and tenofovir disoproxil fumarate tablets may harm your unborn baby. o You should not become pregnant during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Tell you healthcare provider right away if you think you may be pregnant or become pregnant during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.
Females who are able to become pregnant should use effective birth control during treatment with efavirenz, lamivudine and tenofovi  $disoproxil fumarate\ tablets\ and\ for\ 12\ weeks\ after\ stopping\ treatment.\ A\ barrier\ form\ of\ birth\ control\ should\ always\ be\ used\ along\ with$ another type of birth control. If you are able to become pregnant, your healthcare provider should do a pregnancy test before you start efavirenz, lamivudine and

tenofovir disoproxil fumarate tablets. Pregnancy Registry. There is a pregnancy registry for women who take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry.

are breastfeeding or plan to breastfeed. Do not breastfeed if you take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby.Talk to your healthcare provider about the best way to feed your baby.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal Some medicines interact with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Efavirenz, lamivudine and tenofovir disoproxil marate tablets may affect the way other medicines work, and other medicines may affect how efavirenz, lamivudine and tenofovir disoproxil Imarate tablets works. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine. You can ask your healthcare provider or pharmacist for a list of medicines that interact with efavirenz, lamivudine and tenofovir disoproxi fumarate tablets.

Do not start taking a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take efavirenze  $lamivudine\ and\ tenofovir\ disoproxil\ fumarate\ tablets\ with\ other\ medicines.$ low should I take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets' Take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets exactly as your healthcare provider tells you to take it.

Take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets 1 time each day, preferably at bedtime. Taking efavirenz, lamivudine and

enofovir disoproxil fumarate tablets at bedtime might help to make some of the side effects less bothersome Take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets on an empty stomach. Do not miss a dose of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. If you miss a dose, take the missed dose as soon as you remember. If it is almost time for your next dose of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets, do not take the missed dose. Take the next dose at your regular time. Stay under the care of your healthcare provider during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets Do not run out of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. The virus in your blood may increase and the virus may

become harder to treat. When your supply starts to run low, get more from your healthcare provider or pharmacy. If you take too much efavirenz, lamiyudine and tenofoyir disoproxil fumarate tablets, go to the nearest hospital emergency room right away What should I avoid while taking efavirenz, lamivudine and tenofovir disoproxil fumarate tablets?
You should avoid taking medicines that contain sorbitol during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

What are the possible side effects of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets? Ffavirenz Jamiyudine and tenofovir disporoxil furnarate tablets may cause serious side effects including See "What is the most important information I should know about efavirenz, Jamiyudine and tenofovir disonroxil fumarate tablets?" Build-up of an acid in your blood (lactic acidosis). Lactic acidosis can happen in some people who take efavirenz, lamivudine and tenofovi

disoproxil fumarate tablets. Lactic acidosis is a serious medical emergency that can lead to death. Call your healthcare provider right away if you get any of the following symptoms that could be signs of lactic acidosis feel very weak or tired feel cold, especially in your arms and legs unusual (not normal) muscle pain feel dizzy or lightheaded

have a fast or irregular heartbear stomach pain with nausea or vomiting

 $\textbf{Severe liver problems} \ \text{can happen in people who take efavirenz, lamivudine and tenofovir disoproxil furnarate tablets. In some cases, these discontinuous problems in people who take efavirenz, lamivudine and tenofovir disoproxil furnarate tablets. In some cases, these discontinuous problems is a simple of the people who take efavirenz, lamivudine and tenofovir disoproxil furnarate tablets. In some cases, these discontinuous problems is a simple of the people who take efavirenz, lamivudine and tenofovir disoproxil furnarate tablets. In some cases, the second problems is a simple of the people who take efavirenz, lamivudine and tenofovir disoproxil furnarate tablets. In some cases, the second problems is a simple of the people who take efavirenz the people who take efav$ severe liver problems can lead to death. Your liver may become large (hepatomegaly) and you may develop fat in your liver (steatosis) offlammation of your liver (hepatitis) that can lead to liver failure requiring a liver transplant has been reported in some people treated with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Your healthcare provider may do blood tests to check your liver before and during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

weakness

Call your healthcare provider right away if you get any of the following signs or symptoms of liver problems: your skin or the white part of your eyes turns yellow (jaundice) nausea and vomiting dark or "tea-colored" urine pain, aching, or tenderness on the right light-colored stools (bowel movements) side of your stomach-area

stomach (abdomen) swelling You may be more likely to get lactic acidosis or serious liver problems if you are female or very overweight (obese) New or worse kidney problems, including kidney failure. Your healthcare provider may do blood and urine tests to check your kidneys pefore and during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Tell your healthcare provider if you get signs and symptoms of kidney problems, including bone pain that does not go away or worsening bone pain, pain in your arms, hands, legs or feet, broken (fractured) bones, muscle pain or weakness. Serious mental health problems. Get medical help right away if you get any of the following symptoms:

tiredness

are not able to move or speak normal feel sad or hopeless o have thoughts of hurting yourself (suicide) or feel anxious or restless have tried to hurt yourself or others do not trust other people are not able to tell the difference between what is hear or see things that are not real true or real and what is false or unreal Nervous system symptoms are common in people who take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets and can be evere. These symptoms usually begin during the first or second day of treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets and usually go away after 2 to 4 weeks of treatment. Some symptoms may occur months to years after beginning efavirenz,

Symptoms may include: trouble concentrating dizziness trouble sleeping drowsiness o lack of coordination or balance unusual dreams rou have dizziness, trouble concentrating or drowsiness, do not drive a car, use machinery, or do anything that needs you to be alert. Some nervous system symptoms (e.g., confusion, slow thoughts and physical movement, and delusions [false beliefs] or hallucinations [seeing or hearing things that others do not see or hear]) may occur months to years after beginning efavirenz, lamivudine and tenofovir disoproxil

lamivudine and tenofovir disoproxil fumarate tablets therapy. These symptoms may become worse if you drink alcohol, take a medicine for mental health problems, or use certain street drugs during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

marate tablets therapy. Promptly contact your health care provider should any of these symptoms occur. Skin reactions and allergic reactions. Skin reactions or rash can happen and can sometimes be severe. Skin rash usually goes away without any change in treatment. If you develop a rash or a rash with any of the following symptoms, call your healthcare provider right itching peeling skin

swelling of your face red or inflamed eyes blisters or skin lesions

Risk of inflammation of the pancreas (pancreatitis). Children may be at risk for developing pancreatitis during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets if they: have taken nucleoside analogue medicines in the past

have a history of pancreatitis have other risk factors for pancreatitis Call your healthcare provider right away if your child develops signs and symptoms of pancreatitis including severe upper stomach

area pain, with or without nausea and vomiting. Your healthcare provider may tell you to stop giving efavirenz, lamivudine and tenofovir disoproxil fumarate tablets to your child if their symptoms and blood test results show that your child may have pancreatitis. Seizures. Seizures are more likely to happen if you have had seizures in the past. Increases in blood fat levels (cholesterol and triglycerides). Your healthcare provider will check your blood fat levels before and during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

Bone problems can happen in some people who take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Bone problems nclude bone pain, softening or thinning (which may lead to fractures). Your healthcare provider may need to do tests to check your bones. Tell your healthcare provider if you have any bone pain, pain in your hands or feet, or muscle pain or weakness during treatment with

favirenz, lamivudine and tenofovir disoproxil fumarate tablets. Changes in your immune system (Immune Reconstitution Syndrome) can happen when you start taking HIV-1 medicines. Your immune em may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider if you start having new symptoms after starting your HIV-1 medicine. Changes in body fat can happen in some people who take HIV-1 medicines. These changes may include increased amount of fat in the upper back and neck ("buffalo hump"), breast, and around the main part of your body (trunk). Loss of fat from the legs, arms, and face may also

Changes in the electrical activity of your heart called QT prolongation. QT prolongation can cause irregular heartbeats that can be life-threatening. Tell your healthcare provider if you feel faint, lightheaded, dizzy, or feel your heart beating irregularly or fast during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. he most common side effects of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets are rash and dizziness, Fell your healthcare provider if you have any side effect that bothers you or that does not go away

These are not all the possible side effects of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. How should I store efavirenz, lamivudine and tenofovir disoproxil fumarate tablets Store efavirenz, lamivudine and tenofovir disoproxil fumarate tablets at 20° to 25°C (68° to 77°F).

Keep efavirenz, lamivudine and tenofovir disoproxil fumarate tablets and all medicines out of the reach of children. General information about the safe and effective use of efavirenz. Iamivudine and tenofovir disoproxil fumarate tablets. dicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use efavirenz, lamivudine and

tenofovir disoproxil fumarate tablets for a condition for which it was not prescribed. Do not give efavirenz, tamivudine and tenofovir disoproxil fumarate tablets to other people, even if they have the same symptoms that you have. It may harm them. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about efavirenz, lamivudine and tenofovir coproxil fumarate tablets that is written for health professionals. What are the ingredients in efavirenz, lamivudine and tenofovir disoproxil fumarate tablets? Active ingredients: efavirenz USP, lamivudine USP, and tenofovir disoproxil fumarate Inactive ingredients: croscarmellose sodium, ferric oxide vellow, hydroxypropyl cellulose, lactose monohydrate, magnesium stearate crocrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium lauryl sulfate, talc and titanium dioxide

 $Brands\ listed\ are\ the\ trademarks\ of\ their\ respective\ owners\ and\ are\ not\ trademarks\ of\ Laurus\ Labs\ Limited.$ For more information, call Laurus Generics Inc. at 1-833-3-LAURUS (1-833-352-8787) This Patient Information has been approved by the U.S. Food and Drug Administration Manufactured for: Laurus Generics Inc.

Keep efavirenz, lamivudine and tenofovir disoproxil fumarate tablets in the original co

Suite 5200 Berkeley Heights, NJ 07922 Manufactured by Anakapalli-531011 Revised: 08/2023

Revised: 08/2023

Berkeley Heights, NJ 07922 ell Drive

(27 to 50%)

(15 to 40%)

(31 to 58%)

Brands listed are the trademarks of their respective owners and are not trademarks of Laurus Labs Limited For more information, call Laurus Generics Inc. at 1-833-3-LAURUS (1-833-352-8787).

**Inactive ingredients:** croscarmellose sodium, ferric oxide yellow, hydroxypropyl cellulose, lactose monohydrate microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium lauryl sulfate, talc and titanium dioxide.

General information about the safe and effective use of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use efavirenz, la and tenofovir disoproxil fumarate tablets for a condition for which it was not prescribed. Do not give efavirenz, lamivudine and disoproxil fumarate tablets to other people, even if they have the same symptoms that you have. It may harm them. If you would information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about lamivudine and tenofovir disoproxil fumarate tablets that is written for health professionals. What are the ingredients in efavirenz, lamivudine and tenofovir disoproxil fumarate tablets? Active ingredients: efavirenz USP, lamivudine USP, and tenofovir disoproxil fumarate

Keep efavirenz, lamivudine and tenofovir disoproxil fumarate tablets and all medicines out of the reach of children Store efavirenz, lamivudine and tenofovir disoproxil fumarate tablets at 20° to 25°C (68° to 77°F). Keep efavirenz, lamivudine and tenofovir disoproxil fumarate tablets in the original container.

How should I store efavirenz, lamivudine and tenofovir disoproxil fumarate tablets?

Tell your healthcare provider if you have any side effect that bothers you or that does not go mmon side effects of efavirenz, lamivı dine and tenofovir disoproxil fumarate tablets are rash and dizziness.

These are not all the possible side effects of efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. **Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088**.

**Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider if you start having new symptoms after starting your HIV-1 medicine. **Changes in body fat** can happen in some people who take HIV-1 medicines. These changes may include increased amount of fat in the upper back and neck ("buffalo hump"), breast, and around the main part of your body (trunk). Loss of fat from the legs, arms, and face may also happen. The cause and long-term health effects of these conditions are not known. **Changes in the electrical activity of your heart called QT prolongation. QT prolongation can cause irregular heartbeats that can be life-threatening.** Tell your healthcare provider if you feel faint, lightheaded, dizzy, or feel your heart beating irregularly or fast during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

**Bone problems** can happen in some people who take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Bone problems include bone pain, softening or thinning (which may lead to fractures). Your healthcare provider may need to do tests to check your bones. Tell your healthcare provider if you have any bone pain, pain in your hands or feet, or muscle pain or weakness during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Seizures. Seizures are more likely to happen if you have had seizures in the past.

Increases in blood fat levels (cholesterol and triglycerides). Your healthcare provider will check your blood fat levels before and during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets.

Call your healthcare provider right away if your child develops signs and symptoms of pancreatitis including severe upper stomach-area pain, with or without nausea and vomiting. Your healthcare provider may tell you to stop giving efavirenz, lamivudine and tenofovir disoproxil fumarate tablets to your child if their symptoms and blood test results show that your child may have

Risk of inflammation of the pancreas (pancreatitis). Children may be efavirenz, lamivudine and tenofovir disoproxil fumarate tablets if they:

o have taken nucleoside analogue medicines in the past
o have a history of pancreatitis
o have other risk factors for pancreatitis 0 0 0 red or inflamed eyes at risk for developing pancreatitis

swelling of your face blisters or skin lesions

**Skin reactions and allergic reactions.** Skin reactions or rash can happen and can sometimes be severe. Skin rash usually goes away without any change in treatment. If you develop a rash or a rash with any of the following symptoms, call your healthcare provider right

Some nervous system symptoms (e.g., confusion, slow thoughts and physical movement, and delusions [false beliefs] or hallucinations [seeing or hearing things that others do not see or hear]) may occur months to years after beginning efavirenz, lamivudine and tenofovir disoproxil fumarate tablets therapy. Promptly contact your health care provider should any of these symptoms occur. If you have dizziness, trouble concentrating or drowsiness, do not drive a car, use machinery, or do anything that needs you to be alert.

severe. These symptoms are common in people who take efavirenz, lamivudine and tenofovir disoproxil fumarate tablets and can be severe. These symptoms usually begin during the first or second day of treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets and usually go away after 2 to 4 weeks of treatment. Some symptoms may occur months to years after beginning efavirenz, lamivudine and tenofovir disoproxil fumarate tablets therapy. These symptoms may become worse if you drink alcohol, take a medicine for mental health problems, or use certain street drugs during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Symptoms may include:

o dizziness
o trouble sleeping
o drowsiness
o drowsiness

Serious mental health problems. Get medical help right away if you get any of the following symptoms:

o feel sad or hopeless
o feel anxious or restless
o do not trust other people
o hear or see things that are not real
o hear or see things that are not real
o feel anxious or restless
o have tried to hurt yourself or others
o hear or see things that are not real
o are not able to tell the difference between what is have thoughts of hurting yourself (suicide) or have tried to hurt yourself or others are not able to tell the difference between what is true or real and what is false or unreal

You may be more likely to get lactic acidosis or serious liver problems if you are female or very overweight (obese).

• New or worse kidney problems, including kidney failure. Your healthcare provider may do blood and urine tests to check your kidneys before and during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Tell your healthcare provider if you get before and during treatment with efavirenz, lamivudine and tenofovir disoproxil fumarate tablets. Tell your healthcare provider if you get hands, and the favirence including hone pain, pain in your arms, hands, signs and symptoms of kidney problems, including bone pain that does not go away or worsening bone pain, pain in your arms, legs or feet, broken (fractured) bones, muscle pain or weakness. stomach (abdomen) swelling

0 0