

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use GABAPENTIN TABLETS safely and effectively. See full prescribing information for GABAPENTIN TABLETS. GABAPENTIN tablets, for oral use

Initial U.S. Approval: 1993 ----RECENT MAJOR CHANGES--Warnings and Precautions, Respiratory Depression (5.7) 4/2020

----INDICATIONS AND USAGE--Gabapentin tablets are indicated for

• Postherpetic neuralgia in adults (1) Adjunctive therapy in the treatment of partial onset seizures, with and without secondary generalization, in adults and pediatric patients 3 years and older with epilepsy (1)

Postherpetic Neuralgia (2.1)

----DOSAGE AND ADMINISTRATION--

 Dose can be titrated up as needed to a dose of 1,800 mg/day o Day 1: Single 300 mg dose

 $\circ~$ Day 2: 600 mg/day (i.e., 300 mg two times a day)

o Day 3: 900 mg/day (i.e., 300 mg three times a day) • Epilepsy with Partial Onset Seizures (2.2)

Patients 12 years of age and older: starting dose is 300 mg three times daily; may be titrated up to 600 mg three times daily

 Patients 3 to 11 years of age: starting dose range is 10 to 15 mg/kg/day, given in three divided doses; recommended dose in patients 3 to 4 years of age is 40 mg/kg/day, given in three divided doses; the recommended dose in patients 5 to 11 years of age is 25 to 35 mg/kg/day, given in three divided doses. The recommended dose is reached by upward titration over a period of approximately 3 days

• Dose should be adjusted in patients with reduced renal function (2.3, 2.4) -----DOSAGE FORMS AND STRENGTHS----

----CONTRAINDICATIONS-

• Tablets: 600 mg, and 800 mg (3)

Known hypersensitivity to gabapentin or its ingredients (4)

FULL PRESCRIBING INFORMATION: CONTENTS*

INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.2 Dosage for Epilepsy with Partial Onset Seizures

Dosage for Postherpetic Neuralgia

2.3 Dosage Adjustment in Patients with Renal Impairment 2.4 Dosage in Elderly 2.5 Administration Information

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS 5 WARNINGS AND PRECAUTIONS

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan 5.1

Hypersensitivity

Anaphylaxis and Angioedema 5.3

Effects on Driving and Operating Heavy Machinery Somnolence/Sedation and Dizziness 5.4

Withdrawal Precipitated Seizure, Status Epilepticus 5.5 Suicidal Behavior and Ideation

5.7 Neuropsychiatric Adverse Reactions (Pediatric Patients 3 to 12 Years of Age)

5.8

5.9 Tumorigenic Potential 5.10 Sudden and Unexplained Death in Patients with Epilepsy

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

FULL PRESCRIBING INFORMATION

2 DOSAGE AND ADMINISTRATION

2.1 Dosage for Postherpetic Neuralgia

doses should not exceed 12 hours.

1 INDICATIONS AND USAGE Gabapentin Tablets are indicated for:

· Management of postherpetic neuralgia in adults Adjunctive therapy in the treatment of partial onset seizures, with and without secondary generalization, in adults and pediatric patients 3 years and older with epilepsy

In adults with postherpetic neuralgia, gabapentin tablets may be initiated on Day 1 as a single 300 mg dose, on Day 2 as 600 mg/day (300 mg two times a day), and on Day 3 as 900 mg/day (300 mg three times a day). The dose can subsequently be titrated up as needed for pain relief to a dose of 1,800 mg/day (600 mg three times a day). In clinical studies, efficacy was demonstrated over a range of doses from 1,800 mg/day to 3,600 mg/day with comparable effects across the dose range; however, in these clinical studies, the additional benefit of using doses greater than 1,800 mg/day was not demonstrated.

2.2 Dosage for Epilepsy with Partial Onset Seizures Patients 12 years of age and above The starting dose is 300 mg three times a day. The recommended maintenance dose of gabapenting tablets is 300 mg to 600 mg three times a day. Dosages up to 2,400 mg/day have been well tolerated in long-term clinical studies. Doses of 3,600 mg/day have also been administered to a small number of patients for a relatively short duration, and have been well tolerated. Administer gabapentin tablets three times a day using 600 mg or 800 mg tablets. The maximum time between

Pediatric Patients Age 3 to 11 years The starting dose range is 10 mg/kg/day to 15 mg/kg/day, given in three divided doses, and the recommended maintenance dose reached by upward titration over a period of approximately 3 days. The recommended maintenance dose of gabapentin tablets in patients 3 to 4 years of age is 40 mg/kg/day, given in three divided doses. The recommended maintenance dose of gabapentin tablets in patients 5 to 11 years of age is 25 mg/kg/day to 35 mg/kg/day, given in three divided doses. Gabapentin may be administered as the oral solution, capsule, or tablet, or using combinations of these formulations. Dosages up to 50 mg/kg/day have been well tolerated in a

Dosage adjustment in patients 12 years of age and older with renal impairment or undergoing hemodialysis is recommended, as follows (see dosing recommendations above for effective

doses in each indication): TABLE 1. Gabapentin Tablets Dosage Based on Renal Function

Renal Function Creatinine Clearance (mL/min)	Total Daily Dose Range (mg/day)	Dose Regir (mg)	men			
≥ 60	900 to 3,600	300 TID	400 TID	600 TID	800 TID	1,200 TID
>30 to 59	400 to 1,400	200 BID	$300 \; \mathrm{BID}$	400 BID	500 BID	700 BID
>15 to 29	200 to 700	200 QD	300 QD	400 QD	500 QD	700 QD
15a	100 to 300	100 QD	125 QD	150 QD	200 QD	300 QD
	Post-H	Iemodialysis	Supplement	al Dose (mg) ^b	
Hemodialysis		125 ^b	150 ^b	200 ^b	250b	350 ^b

TID = Three times a day; BID = Two times a day; QD = Single daily dose

For patients with creatinine clearance <15 mL/min, reduce daily dose in proportion to creatinine clearance (e.g., patients with a creatinine clearance of 7.5 mL/min should receive one-half the daily dose that patients with a creatinine clearance of 15 mL/min receive).

Patients on hemodialysis should receive maintenance doses based on estimates of creatining clearance as indicated in the upper portion of the table and a supplemental post-hemodialysis dose administered after each 4 hours of hemodialysis as indicated in the lower portion of the Creatinine clearance (CLCr) is difficult to measure in outpatients. In patients with stable renal

function, creatinine clearance can be reasonably well estimated using the equation of Cockcroft [140 - age (years)] x weight (kg)

CLCr = (x 0.85 for female patients) 72 x serum creatinine (mg/dL)

The use of gabapentin tablets in patients less than 12 years of age with compromised renal function has not been studied.

Because elderly patients are more likely to have decreased renal function, care should be taken

2.5 Administration Information

Administer gabapentin tablets orally with or without food.

Inform patients that, should they divide the scored 600 mg or 800 mg gabapentin tablets in order to administer a half-tablet, they should take the unused half-tablet as the next dose. Half-tablets not used within 28 days of dividing the scored tablet should be discarded.

needed at the discretion of the prescriber).

If the gabapentin tablets dose is reduced, discontinued, or substituted with an alternative on, this should be done gradually over a minimum of 1 week (a longer period may be

3 DOSAGE FORMS AND STRENGTHS **Tablets**

• 600 mg: White to off white, elliptical, biconvex film-coated tablets functionally scored on both sides, debossed with "C" and "5" on either side of score line on one side.

• 800 mg: White to off white, elliptical, biconvex film-coated tablets functionally scored or both sides, debossed with "C" and "7" on either side of score line on one side. 4 CONTRAINDICATIONS

Gabapentin tablets are contraindicated in patients who have demonstrated hypersensitivity to

5 WARNINGS AND PRECAUTIONS 5.1 Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS), also known as multiorgan hypersensitivity, has occurred with gabapentin. Some of these reactions have been fatal or

life-threatening. DRESS typically, although not exclusively, presents with fever, rash, and/or lymphadenopathy, in association with other organ system involvement, such as hepatitis, nephritis, hematological abnormalities, myocarditis, or myositis sometimes resembling an acute viral infection. Eosinophilia is often present. This disorder is variable in its expression, and other organ systems not noted here may be involved.

It is important to note that early manifestations of hypersensitivity, such as fever or lymphadenopathy, may be present even though rash is not evident. If such signs or symptoms are

new,

are

if they

especially

if you have any of these

Call a healthcare provider right

de

 attempts to commit suicie new or worse depression

• thoughts about suicide or

worse, or worry you:

o

suicidal thoughts

, gabapentin t in 500.

r antiepileptic drugs, g er of people, about 1 in t away if you have an

number

actions in a very small

1. Suicidal Thoughts. Like other

Gabapentin tablets can cause

Stopping gabapentin tablet

including:

effects

serious side can

healthcare provider.

to your

ts without first

cause

pentin tablets ts suddenly c

Do not stop taking gabapentin

What is the most importa

(gab"

----WARNINGS AND PRECAUTIONS--

• Drug Reaction with Eosinophilia and Systemic Symptoms (Multiorgan hypersensitivity): Discontinue if alternative etiology is not established (5.1)

• Anaphylaxis and Angioedema: Discontinue and evaluate patient immediately (5.2) Driving Impairment: Somnolence/Sedation and Dizziness: Warn patients not to drive until

they have gained sufficient experience to assess whether their ability to drive or operate heavy machinery will be impaired (5.3, 5.4) Increased seizure frequency may occur in patients with seizure disorders if gabapentin is

abruptly discontinued (5.5) Suicidal Behavior and Ideation: Monitor for suicidal thoughts/behavior (5.6)

· Respiratory Depression: May occur with gabapentin when used with concomitant central

respiratory Depression. May occur with gaoapenin when used with concominant central nervous system (CNS) depressants, including opioids, or in the setting of underlying respiratory impairment. Monitor patients and adjust dosage as appropriate (5.7) Neuropsychiatric Adverse Reactions in Children 3 to 12 Years of Age: Monitor for such

----ADVERSE REACTIONS-Most common adverse reactions (incidence \geq 8% and at least twice that for placebo) were: • Postherpetic neuralgia: Dizziness, somnolence, and peripheral edema (6.1) • Epilepsy in patients >12 years of age: Somnolence, dizziness, ataxia, fatigue, and nystagmus

• Epilepsy in patients 3 to 12 years of age: Viral infection, fever, nausea and/or vomiting, somnolence, and hostility (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Laurus Generics Inc. at 1-833-3-LAURUS (1-833-352-8787) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. -----DRUG INTERACTIONS--

Concentrations increased by morphine; may need dose adjustment (5.4, 7.1) ----USE IN SPECIFIC POPULATIONS

Pregnancy: Based on animal data, may cause fetal harm (8.1) See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

Revised: 10/2023

7.2 Other Antiepileptic Drugs

Maalox® (aluminum hydroxide, magnesium hydroxide) 7.4 Drug/Laboratory Test Interactions

8 USE IN SPECIFIC POPULATIONS 8.1 Pregnancy

8.2 Lactation

events (5.8)

Pediatric Use

Geriatric Use 8.6 Renal Impairme 9 DRUG ARUSE AND DEPENDENCE

Controlled Substance Abuse

9.3 Dependence 10 OVERDOSAGE

11 DESCRIPTION 12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES 14.1 Postherpetic Neuralgia

14.2 Epilepsy for Partial Onset Seizures (Adjunctive Therapy)

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION *Sections or subsections omitted from the full prescribing information are not listed.

present, the patient should be evaluated immediately. Gabapentin should be discontinued if an

alternative etiology for the signs or symptoms cannot be established.

5.2 Anaphylaxis and Angioedema Gabapentin can cause anaphylaxis and angioedema after the first dose or at any time during treatment. Signs and symptoms in reported cases have included difficulty breathing, swelling of

the lips, throat, and tongue, and hypotension requiring emergency treatment. Patients should be instructed to discontinue gabapentin and seek immediate medical care should they experience signs or symptoms of anaphylaxis or angioedema.

5.3 Effects on Driving and Operating Heavy Machinery Patients taking gabapentin should not drive until they have gained sufficient experience to assess whether gabapentin impairs their ability to drive. Driving performance studies conducted with a prodrug of gabapentin (gabapentin enacarbil tablet, extended-release) indicate that gabapentin may cause significant driving impairment. Prescribers and patients should be aware that patients' ability to assess their own driving competence, as well as their ability to assess the degree of somnolence caused by gabapentin, can be imperfect. The duration of driving impairment after starting therapy with gabapentin is unknown. Whether the impairment is related to somnolence

[see Warnings and Precautions (5.4)] or other effects of gabapentin is unknown. Moreover, because gabapentin causes somnolence and dizziness [see Warnings and Precautions (5.4)], patients should be advised not to operate complex machinery until they have gained sufficient experience on gabapentin to assess whether gabapentin impairs their ability to perform

such tasks.

5.4 Somnolence/Sedation and Dizziness During the controlled epilepsy trials in patients older than 12 years of age receiving doses of gabapentin up to 1,800 mg daily, somnolence, dizziness, and ataxia were reported at a greater rate in patients receiving gabapentin compared to placebo: i.e., 19% in drug versus 9% in placebo for somnolence, 17% in drug versus 7% in placebo for dizziness, and 13% in drug versus 6% in placebo for ataxia. In these trials somnolence, ataxia and fatigue were common adverse reactions leading to discontinuation of gabapentin in patients older than 12 years of age, with 1.2%, 0.8% and 0.6% discontinuing for these events, respectively.

During the controlled trials in patients with post-herpetic neuralgia, somnolence, and dizziness were reported at a greater rate compared to placebo in patients receiving gabapentin, in dosages up to 3,600 mg per day: i.e., 21% in gabapentin-treated patients versus 5% in placebo-treated patients for somnolence and 28% in gabapentin-treated patients versus 8% in placebo-treated patients for somnolence and 28% in gabapentin-treated patients versus 8% in placebo-treated patients for dizziness. Dizziness and somnolence were among the most common adverse reactions leading to discontinuation of gabapentin.

Patients should be carefully observed for signs of central nervous system (CNS) depression, such as sommolence and sedation, when gabapentin is used with other drugs with sedative properties because of potential synergy. In addition, patients who require concomitant treatment with morphine may experience increases in gabapentin concentrations and may require dose adjustment [see Drug Interactions (7.1)].

5.5 Withdrawal Precipitated Seizure, Status Epilepticus Antiepileptic drugs should not be abruptly discontinued because of the possibility of increasing

In the placebo-controlled epilepsy studies in patients >12 years of age, the incidence of status epilepticus in patients receiving gabapentin was 0.6% (3 of 543) versus 0.5% in patients receiving placebo (2 of 378). Among the 2,074 patients >12 years of age treated with gabapentin across all epilepsy studies (controlled and uncontrolled), 31 (1.5%) had status epilepticus. Of these, 14 patients had no prior history of status epilepticus either before treatment or while on other medications. Because adequate historical data are not available, it is impossible to say whether or not treatment with gabapentin is associated with a higher or lower rate of status epilepticus than would be expected to occur in a similar population not treated with gabapentin.

Antiepileptic drugs (AEDs), including gabapentin, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% Cl:1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27.863 AED-treated patients was 0.43%, compared to 0.24% among 16,029 placebo-treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients an increase of approximately one case of such an immig of behavior of every 550 pt treated. There were four suicides in drug-treated patients in the trials and none in placebo-patients, but the number is too small to allow any conclusion about drug effect on suicide. The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5 to 100 years) in the clinical trials analyzed. Table 2 shows absolute and relative risk by indication for all evaluated AEDs.

thoughts or behavior beyond 24 weeks could not be assessed.

TABLE 2 Risk by Indication for Antiepileptic Drugs in the Pooled Analysis Placebo Patients Drug Patients Relative Risk: Risk Difference

indication	with Events Per 1,000 Patients	with Events Per 1,000 Patients	Incidence of Events in Drug Patients/ Incidence in Placebo Patients	Additional Drug Patients with Events Per 1,000 Patients
Epilepsy	1.0	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9
	k for suicidal thought trials for psychiatric			

similar for the epilepsy and psychiatric indications.

Anyone considering prescribing gabapentin or any other AED must balance the risk of suicidal thoughts or behavior with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated. Patients, their caregivers, and families should be informed that AEDs increase the risk of

suicidal thoughts and behavior and should be advised of the need to be alert for the emergence or worsening of the signs and symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts about self-harm. Behaviors of 5.7 Respiratory Depression

There is evidence from case reports, human studies, and animal studies associating gabapentin

Ξ.

medicine suddenly

seizure 1

can cause serious problems.

Do not stop taking gabapentin tablets
Stopping gabapentin tablets suddenly a patient who has epilepsy can cause s

seizures that will not stop

healthca

æ

t0

talking

tablets without first

especially

as

visits

. Stopping a seizure status epilepticus).

behaviors, thoughts,

especially sudden changes, in mood,

symptoms

ıges,

Pay attention to any chan

How can I watch for earl

(mania)

an extreme increase in activity

other unusual changes in

acting on dangerous impulses

• trouble sleeping (insomnia)

 new or worse irritability acting aggressive, being

• feeling agitated or restless

• panic attacks

new or worse anxiety

scheduled.

provider as

your 1

with your

• Keep all follow-up visits 'Call your healthcare provider

Ö

suicidal thoughts

caused by things other than medicines. ay check for other causes.

ons can be provider

• Suicidal thoughts or actions actions, your healthcare pro-

cause school

can in s

ars of age changes

children 3 to 12 years ation, restlessness, ch

concentration,

- Using gabapentin tablets ior, problems with concer

with serious, life-threatening, or fatal respiratory depression when coadministered with CNS depressants, including opioids, or in the setting of underlying respiratory impairment. When the decision is made to co-prescribe gabapentin with another CNS depressant, particularly an opioid, or to prescribe gabapentin to patients with underlying respiratory impairment, monitor patients for symptoms of respiratory depression and sedation, and consider initiating gabapentin at a low dose. The management of respiratory depression may include close observation, supportive

ures, and reduction or withdrawal of CNS depressants (including gabapentin 5.8 Neuropsychiatric Adverse Reactions (Pediatric Patients 3 to 12 Years of Age)

Gabapentin use in pediatric patients with epilepsy 3 to 12 years of age is associated with the occurrence of CNS related adverse reactions. The most significant of these can be classified into the following categories: 1) emotional lability (primarily behavioral problems), 2) hostility, including aggressive behaviors, 3) thought disorder, including concentration problems and change in school performance, and 4) hyperkinesia (primarily restlessness and hyperactivity).

Among the gabapentin-treated patients, most of the reactions were mild to moderate in intensity. In controlled clinical epilepsy trials in pediatric patients 3 to 12 years of age, the incidence of these adverse reactions was: emotional lability 6% (gabapentin-treated patients) versus 1.3% (placebo-freated patients); hostility 5.2% versus 1.3%; hyperkinesia 4.7% versus 2.9%; and thought disorder 1.7% versus 0%. One of these reactions, a report of hostility, was considered serious. Discontinuation of gabapentin treatment occurred in 1.3% of patients reporting emotional lability and hyperkinesia and 0.9% of gabapentin-treated patients reporting hostility and thought disorder. One placebo-treated patient (0.4%) withdrew due to emotional lability.

5.9 Tumorigenic Potential

In an oral carcinogenicity study, gabapentin increased the incidence of pancreatic acinar cell tumors in rats [see Nonclinical Toxicology (13.1)]. The clinical significance of this finding is unknown. Clinical experience during gabapentin's premarketing development provides no direct means to assess its potential for inducing tumors in humans.

In clinical studies in adjunctive therapy in epilepsy comprising 2,085 patient-years of exposure in patients >12 years of age, new tumors were reported in 10 patients (2 breast, 3 brain, 2 lung, 1 adrenal, 1 non-Hodgkin's lymphoma, 1 endometrial carcinoma *in situ*), and preexisting tumors worsened in 11 patients (9 brain, 1 breast, 1 prostate) during or up to 2 years following discontinuation of gabapentin. Without knowledge of the background incidence and recurrent in a similar population not treated with gabapentin, it is impossible to know whether the incidence seen in this cohort is or is not affected by treatment.

5.10 Sudden and Unexplained Death in Patients with Enilepsy

During the course of premarketing development of gabapentin, 8 sudden and unexplained deaths were recorded among a cohort of 2,203 epilepsy patients treated (2,103 patient-years of exposure) with gabapentin

Some of these could represent seizure-related deaths in which the seizure was not observed, e.g., at night. This represents an incidence of 0.0038 deaths per patient-year. Although this rate exceeds that expected in a healthy population matched for age and sex, it is within the range of estimates for the incidence of sudden unexplained deaths in patients with epilepsy not receiving gabapentin (ranging from 0.0005 for the general population of epileptics to 0.003 for a clinical trial population similar to that in the gabapentin program, to 0.005 for patients with refractory epilepsy). Consequently, whether these figures are reassuring or raise further concern depends a comparability of the populations reported upon to the gabapentin cohort and the accuracy of

6 ADVERSE REACTIONS The following serious adverse reactions are discussed in greater detail in other sections:

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity [see Warnings and Precautions (5.1)]

Anaphylaxis and Angioedema [see Warnings and Precautions (5.2)]

• Somnolence/Sedation and Dizziness [see Warnings and Precautions (5.4)] • Withdrawal Precipitated Seizure, Status Epilepticus [see Warnings and Precautions (5.5)] Suicidal Behavior and Ideation [see Warnings and Precautions (5.6)]

 Respiratory Depression [see Warnings and Precautions (5.7)]. • Neuropsychiatric Adverse Reactions (Pediatric Patients 3 to 12 Years of Age) [see Warnings

and Precautions (5.8)] · Sudden and Unexplained Death in Patients with Epilepsy [see Warnings and Precautions

6.1 Clinical Trials Experience Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Postherpetic Neuralgia The most common adverse reactions associated with the use of gabapentin in adults, not seen at an equivalent frequency among placebo-treated patients, were dizziness, somnolence, and peripheral edema

In the 2 controlled trials in postherpetic neuralgia, 16% of the 336 patients who received gabapentin and 9% of the 227 patients who received placebo discontinued treatment because of an adverse reaction. The adverse reactions that most frequently led to withdrawal in gabapentintreated patients were dizziness, somnolence, and nausea. Table 3 lists adverse reactions that occurred in at least 1% of gabapentin-treated patients with

postherpetic neuralgia participating in placebo-controlled trials and that were numerically more frequent in the gabapentin group than in the placebo group. TABLE 3. Adverse Reactions in Pooled Placebo-Controlled Trials in Postherpetic Neuralgia

	Gabapentin N=336	Placebo N=227
	%	%
Body as a Whole		
Asthenia	6	5
Infection	5	4
Accidental injury	3	1
Digestive System		
Diarrhea	6	3
Dry mouth	5	1
Constipation	4	2
Nausea	4	3
Vomiting	3	2
Metabolic and Nutritional Disorders		
Peripheral edema	8	2
Weight gain	2	0
Hyperglycemia	1	0
Nervous System		
Dizziness	28	8
Somnolence	21	5
Ataxia	3	0
Abnormal thinking	3	0
Abnormal gait	2	0
Incoordination	2	0
Respiratory System		
Pharyngitis	1	0
Special Senses		
Amblyopia ^a	3	1
Conjunctivitis	1	0
Diplopia	1	0

Otitis media Reported as blurred vision Other reactions in more than 1% of patients but equally or more frequent in the placebo group

[see Warnings and Precautions (5.8)].

Urogenital System

nay affect your skin ospitalized or to stop gabapentin tablets.

This may a allergic r

blood cells. The rash with an a

There were no clinically important differences between men and women in the types incidence of adverse reactions. Because there were few patients whose race was reported as other than white, there are insufficient data to support a statement regarding the distribution of adverse reactions by race. Epilepsy with Partial Onset Seizures (Adjunctive Therapy)

included pain, tremor, neuralgia, back pain, dyspepsia, dyspnea, and flu syndrome

The most common adverse reactions with gabapentin in combination with other antiepileptic drugs in patients >12 years of age, not seen at an equivalent frequency among placebo-treated patients, were somnolence, dizziness, ataxia, fatigue, and nystagmus. The most common adverse reactions with gabapentin in combination with other antiepileptic drugs in pediatric patients 3 to 12 years of age, not seen at an equal frequency among placebo treated patients, were viral infection, fever, nausea and/or vomiting, somnolence, and hostility

liatric patients 3 to 12 years of age who received gabapentin in premarketing clinical trials discontinued treatment because of an adverse reaction. The adverse reactions most commonly associated with withdrawal in patients >12 years of age were somnolence (1.2%), ataxis (0.8%), fatigue (0.6%), nausea and/or vomiting (0.6%), and dizziness (0.6%). The adverse reactions most commonly associated with withdrawal in pediatric patients were emotional lability (1.6%), hostility (1.3%), and hyperkinesia (1.1%). Table 4 lists adverse reactions that occurred in at least 1% of gabapentin-treated patients >12

Approximately 7% of the 2,074 patients >12 years of age and approximately 7% of the 449

years of age with epilepsy participating in placebo-controlled trials and were numerically more common in the gabapentin group. In these studies, either gabapentin or placebo was added to the patient's current antiepileptic drug therapy. TABLE 4. Adverse Reactions in Pooled Placebo-Controlled Add-On Trials In Epilepsy

N=543

N = 378

starting

s taken by so hing when s

gabape،.. r when it is taken مراهم المالية add breathing when

awareness, or whe

or

breathing

	%	%
Body As A Whole		
Fatigue	11	5
Increased Weight	3	2
Back Pain	2	1
Peripheral Edema	2	1
Cardiovascular		
Vasodilatation	1	0
Digestive System		
Dyspepsia	2	1
Dry Mouth or Throat	2	1
Constipation	2	1
Dental Abnormalities	2	0
Nervous System		
Somnolence	19	9
Dizziness	17	7
Ataxia	13	6
Nystagmus	8	4
Tremor	7	3
Dysarthria	2	1
Amnesia	2	0
Depression	2	1
Abnormal thinking	2	1
Abnormal coordination	1	0
Respiratory System		
Pharyngitis	3	2
Coughing	2	1
Skin and Appendages		
Abrasion	1	0

Special Senses Amblyopia^b

a Plus background antiepileptic drug therapy

b Amblyopia was often described as blurred vision. Among the adverse reactions occurring at an incidence of at least 10% in gabapentin-treated

patients, somnolence and ataxia appeared to exhibit a positive dose-response relationship. The overall incidence of adverse reactions and the types of adverse reactions seen were similar among men and women treated with gabapentin. The incidence of adverse reactions increased slightly with increasing age in patients treated with either gabapentin or placebo. Because only 3% of patients (28/921) in placebo-controlled studies were identified as nonwhite (black or other), there are insufficient data to support a statement regarding the distribution of adverse

Table 5 lists adverse reactions that occurred in at least 2% of gabapentin-treated patients, age 3 to 12 years of age with epilepsy participating in placebo-controlled trials, and which were numerically more common in the gabanentin group

TABLE 5. Adverse Reactions in a Placebo-Controlled Add-On Trial in Pediatric Epilepsy

	Gabapentin ^a N=119 %	Placebo ^a N=128
Body As A Whole		
Viral Infection	11	3
Fever	10	3
Increased Weight	3	1
Fatigue	3	2
Digestive System		
Nausea and/or Vomiting	8	7
Nervous System		
Somnolence	8	5
Hostility	8	2
Emotional Lability	4	2
Dizziness	3	2
Hyperkinesia	3	1
Respiratory System		
Bronchitis	3	1
Respiratory Infection	3	1

a Plus background antiepileptic drug therapy

Other reactions in more than 2% of pediatric patients 3 to 12 years of age but equally or more frequent in the placebo group included: pharyngitis, upper respiratory infection, headache, rhinitis, convulsions, diarrhea, anorexia, coughing, and otitis media 6.2 Postmarketing Experience

The following adverse reactions have been identified during postmarketing use of gabapentin

7.2 Other Antiepileptic Drugs

icology (12.3)].

8.1 Pregnancy

[see Data].

8 USE IN SPECIFIC POPULATIONS

Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug

Hepatobiliary disorders: jaundice

Investigations: elevated creatine kinase, elevated liver function tests Metabolism and nutrition disorders: hyponatremia

Musculoskeletal and connective tissue disorder: rhabdomyolysis Nervous system disorders: movement disorder

Psychiatric disorders: agitation Reproductive system and breast disorders: breast enlargement, changes in libido, ejaculation disorders and anorgasmia

Skin and subcutaneous tissue disorders: angioedema [see Warnings and Precautions (5.2)], bullous pemphigoid, erythema multiforme, Stevens-Johnson syndrome, There are postmarketing reports of life-threatening or fatal respiratory depression in patients taking gabapentin with opioids or other CNS depressants, or in the setting of underlying respiratory impairment [see Warnings and Precautions (5.7)].

Adverse reactions following the abrupt discontinuation of gabapentin have also been reported. The most frequently reported reactions were anxiety, insomnia, nausea, pain, and sweating 7.1 Opioids Respiratory depression and sedation, sometimes resulting in death, have been reported following

buprenorphine) [see Warnings and Precautions (5.7)]. Hydrocodone Coadministration of gabapentin with hydrocodone decreases hydrocodone exposure [see Clinical Pharmacology (12.3)]. The potential for alteration in hydrocodone exposure and effect should be considered when gabapentin is started or discontinued in a patient taking hydrocodone exposure.

coadministration of gabapentin with opioids (e.g., morphine, hydrocodone, oxycodone,

Morphine When gabapentin is administered with morphine, patients should be observed for signs of CNS depression, such as somnolence, sedation and respiratory depression [see Clinical Pharm (12.3)1.

commonly coadministered antiepileptic drugs [see Clinical Pharmacology (12.3)]. 7.3 Maalox® (aluminum hydroxide, magnesium hydroxide) The mean bioavailability of gabapentin was reduced by about 20% with concomitant use of an antacid (Maalox®) containing magnesium and aluminum hydroxides. It is recommended that gabapentin be taken at least 2 hours following Maalox administration [see Clinical

Gabapentin is not appreciably metabolized nor does it interfere with the metabolism of

7.4 Drug/Laboratory Test Interactions Because false positive readings were reported with the Ames N-Multistix SG® dipstick test for urinary protein when gabapentin was added to other antiepilentic drugs, the more specific sulfosalicylic acid precipitation procedure is recommended to determine the presence of urine

 $There \ are \ no \ adequate \ data \ on \ the \ developmental \ risks \ associated \ with \ the \ use \ of \ gabapentin \ in pregnant \ women. \ In \ nonclinical \ studies \ in \ mice, \ rats, \ and \ rabbits, \ gabapentin \ was \ developmentally$ toxic (increased fetal skeletal and visceral abnormalities, and increased embryofetal mortality) when administered to pregnant animals at doses similar to or lower than those used clinically

In the U.S. general population, the estimated background risk of major birth defects and

miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively. The background risk of major birth defects and miscarriage for the indicated population is unknown. Animal data When pregnant mice received oral doses of gabanentin (500, 1,000, or 3,000 mg/kg/day) during the period of organogenesis, embryofetal toxicity (increased incidences of skeletal variations) was observed at the two highest doses. The no-effect dose for embryofetal developmental

toxicity in mice (500 mg/kg/day) is less than the maximum recommended human dose (MRHD) of 3,600 mg on a body surface area (mg/m²) basis. In studies in which rats received oral doses of gabapentin (500 to 2,000 mg/kg/day) during pregnancy, adverse effect on offspring development (increased incidences of hydroureter and

hydronephrosis) were observed at all doses. The lowest dose tested is similar to the MRHD

When pregnant rabbits were treated with gabapentin during the period of organoger increase in embryofetal mortality was observed at all doses tested (60, 300, or 1.500 mg/kg). The lowest dose tested is less than the MRHD on a mg/m² basis. In a published study, gabapentin (400 mg/kg/day) was administered by intraperitoneal injection to neonatal mice during the first postnatal week, a period of synaptogenesis in rodents (corresponding to the last trimester of pregnancy in humans). Gabapentin caused a marked decrease in neuronal synapse formation in brains of intact mice and abnormal neuronal synapse

formation in a mouse model of synaptic repair. Gabapentin has been shown in vitro to interfere

with activity of the a28 subunit of voltage-activated calcium channels, a receptor involved in neuronal synaptogenesis. The clinical significance of these findings is unknown. Risk Summary Gabapentin is secreted in human milk following oral administration. The effects on the breastfed infant and on milk production are unknown. The developmental and health benefits

on a mg/m2 basis.

of breastfeeding should be considered along with the mother's clinical need for gabapentin and any potential adverse effects on the breastfed infant from gabapentin or from the underlying maternal condition. 8.4 Pediatric Use

Safety and effectiveness of gabapentin in the management of postherpetic neuralgia in pediatric

Safety and effectiveness as adjunctive therapy in the treatment of partial seizures in pediatric

were 75 years of age and older. There was a larger treatment effect in patients 75 years of age and

patients below the age of 3 years has not been established [see Clinical Studies (14.2)]. 8.5 Geriatric Use The total number of patients treated with gabapentin in controlled clinical trials in patients with postherpetic neuralgia was 336, of which 102 (30%) were 65 to 74 years of age, and 168 (50%)

older compared to younger patients who received the same dosage. Since gabapentin is almost exclusively eliminated by renal excretion, the larger treatment effect observed in patients ≥75 years may be a consequence of increased gabapentin exposure for a given dose that results from an age-related decrease in renal function. However, other factors cannot be excluded. The types and incidence of adverse reactions were similar across age groups except for peripheral edema and ataxia, which tended to increase in incidence with age Clinical studies of gabapentin in epilepsy did not include sufficient numbers of subjects aged 65 and over to determine whether they responded differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger

the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy. This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions are more likely to have decreased renal function, care should be taken in dose selection, and dose should be adjusted based on creatinine clearance values in these patients [see Dosage and Administration (2.4), Adverse Reactions (6), and Clinical Pharmacology (12.3)].

Dosage adjustment in adult patients with compromised renal function is necessary [see Dosage

and Administration (2.3) and Clinical Pharmacology (12.3)]. Pediatric patients with renal

patients. In general, dose selection for an elderly patient should be cautious, usually starting at

Dosage adjustment in patients undergoing hemodialysis is necessary [see Dosage and Administration (2.3) and Clinical Pharmacology (12.3)].

9.1 Controlled Substance

 $_{\text{jo}}$

children

ii.

comes

rash that

painful 1

shingles (a)

 $\quad \text{of} \quad$

8.6 Renal Impairment

Abuse is the intentional, non-therapeutic use of a drug, even once, for its desirable psychological or physiological effects. Misuse is the intentional use, for therapeutic purposes, of a drug by

9 DRUG ABUSE AND DEPENDENCE Gabapentin is not a scheduled drug

gabapentin

ts in garablets

gabapentin

ingredients in gabapentin tablets?

of

of

provider if you:

behavior

suicidal thoughts or

or

problems,

hemodialysis

on

known if gabapentin tablets can harm your unborn baby. scome pregnant while taking gabapentin tablets. You and I take gabapentin tablets while you are pregnant. So an into breast milk. You and your healthcare on the pass into breast milk. breast milk. gabapentin t into take t is not known Eyou become p a should take g Gabapentin o

away if y if you s me pregnant. I r right away if I decide if you breast-feed. (are pregnant or plan to become ider ider will o to your healthcare prov are breast-feeding or plai provider should decide ho your healthcare provider

emotional changes, aggressive behavior, provider may chec emotional changes, aggressive behavior, programance, and hyperactivity.

3. Gabapentin tablete

3. Gabapentin tablets may cause or other parts of your body such as gabapentin tablets. You may or m: Call a healthcare provider right aw

 skin rash hives

• swelling of your face, lips, throat, swollen glands that do not yellowing of your skin or

of the whites of the eyes

tongue

g

difficulty breathing

fever

unusual bruising or bleeding severe fatigue or weakness

 unexpected muscle pain frequent infections

tablets. gabapentin Serions signs ing gab taki These symptoms may be the decide if you should continue

A

vere sleepiness or decreased Watch for increased sleepir problems

4. Serious breathing problems. with other medicines that can

increased Get help ri an cause sevent problems. Very the dose is i lets? What are gabapentin tabl who already has breathing gabapentin tablets or wher

Gabapentin tablets are a prescription Pain from damaged nerv

'es (postherpetic pain) that follows healing ction) in adults. other ection) in adul ken together v Partial seizures when tal after a herpes zoster infe with seizures.

olets if you are allergic to gabapentin Medication Guide for a complete list Who should not take gabapentin Do not take gabapentin tab tablets. See the end of this

What should I tell my healthcare provider before taking. Before taking gabapentin tablets, tell your healthcare pro

poom problems sion, have or have had kidney

 have or have had depres have diabetes

have breathing problems Tell

Front Page

PC:768

an individual in a way other than prescribed by a health care provider or for whom it was not

Gabapentin does not exhibit affinity for benzodiazepine, opioid (mu, delta or kappa), or cannabinoid 1 receptor sites. Gabapentin misuse and abuse have been reported in the postmarketing setting and published literature. Most of the individuals described in these reports had a history of polysubstance abuse. Some of these individuals were taking higher than recommended doses of gabapentin for unapproved uses. When prescribing gabapentin carefully evaluate patients for a history of drug abuse and observe them for signs and symptoms of gabapentin misuse or abuse (e.g., self-dose escalation and drug-seeking behavior). The abuse potential of gabapentin has not been evaluated in human studies

9.3 Dependence Physical dependence is a state that develops as a result of physiological adaptation in response to repeated drug use, manifested by withdrawal signs and symptoms after abrupt discont or a significant dose reduction of a drug. There are rare postmarketing reports of individuals experiencing withdrawal symptoms shortly after discontinuing higher than recommended doses of gabapentin used to treat illnesses for which the drug is not approved. Such symptoms included agitation, disorientation and confusion after suddenly discontinuing gabapentin that resolved after restarting gabapentin. The dependence potential of gabapentin has not been evaluated in human studies

10 OVERDOSAGE

Signs of acute toxicity in animals included ataxia, labored breathing, ptosis, sedation, ypoactivity, or excitation

Acute oral overdoses of gabapentin have been reported. Symptoms have included double vision tremor, slurred speech, drowsiness, altered mental status, dizziness, lethargy, and diarrhea. Fatal respiratory depression has been reported with gabapentin overdose, alone and in combination with other CNS depressants Gabapentin can be removed by hemodialysis.

If overexposure occurs, call your poison control center at 1-800-222-1222

11 DESCRIPTION

The active ingredient in gabapentin tablets, USP is gabapentin USP, which has the chemical name 1-(Aminomethyl)cyclohexaneacetic acid. The molecular formula of gabapentin is C_aH₁₂NO₂ and the molecular weight is 171.24. The

structural formula of gabapentin is -соон

Gabapentin USP is a white to off-white crystalline solid. It is freely soluble in water and in alkaline and acidic solutions. The log of the partition coefficient (n-octanol/0.05M phosphate buffer) at pH 7.4 is -1.25.

Each gabapentin tablet, USP contains 600 mg or 800 mg of gabapentin USP and the following nactive ingredients: copovidone, maize starch 5%, microcrystalline cellulose, magnesium stearate and poloxamer P407. In addition, the film coating contains the following inactive ingredients: Polyethylene Glycol 3350, Polyvinyl alcohol-part hydrolyzed, titanium oxide and

FDA approved dissolution testing specifications differ from USP.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action The precise mechanisms by which gabapentin produces its analgesic and antiepileptic actions are unknown. Gabapentin is structurally related to the neurotransmitter gamma-aminobutyric acid (GABA) but has no effect on GABA binding, uptake, or degradation. *In vitro* studies have shown that gabapentin binds with high-affinity to the $\alpha 2\delta$ subunit of voltage-activated calcium channels; however, the relationship of this binding to the therapeutic effects of gabapentin is unknown.

12.3 Pharmacokinetics All pharmacological actions following gabapentin administration are due to the activity of the parent compound; gabapentin is not appreciably metabolized in humans

Oral Bioavailability

Gabapentin bioavailability is not dose proportional; i.e., as dose is increased, bioavailability decreases. Bioavailability of gabapentin is approximately 60%, 47%, 34%, 33%, and 27% following 900, 1,200, 2,400, 3,600, and 4,800 mg/day given in 3 divided doses, respectively. Food has only a slight effect on the rate and extent of absorption of gabapentin (14% increase in AUC and C_{max}). Distribution

Less than 3% of gabapentin circulates bound to plasma protein. The apparent volume of distribution of gabapentin after 150 mg intravenous administration is 58±6 L (mean ±SD). In patients with epilepsy, steady-state predose (C_{min}) concentrations of gabapentin in cerebrospinal fluid were approximately 20% of the corresponding plasma concentrations Elimination

Gabapentin is eliminated from the systemic circulation by renal excretion as unchanged drug. Gabapentin is not appreciably metabolized in humans.

Gabapentin elimination half-life is 5 to 7 hours and is unaltered by dose or following multiple dosing. Gabapentin elimination rate constant, plasma clearance, and renal clearance are directly proportional to creatinine clearance. In elderly patients, and in patients with impaired renal function, gabapentin plasma clearance is reduced. Gabapentin can be removed from plasma by hemodialysis. Specific Populations

The effect of age was studied in subjects 20 to 80 years of age. Apparent oral clearance (CL/F) of gabapentin decreased as age increased, from about 225 mL/min in those under 30 years of age to about 125 mL/min in those over 70 years of age. Renal clearance (CLr) and CLr adjusted for body surface area also declined with age; however, the decline in the renal clearance of gabapentin with age can largely be explained by the decline in renal function. I see Dosage and Administration (2.4) and Use in Specific Populations (8.5)]. Gender

Although no formal study has been conducted to compare the pharmacokinetics of gabapentin in men and women, it appears that the pharmacokinetic parameters for males and females are similar and there are no significant gender differences.

Pharmacokinetic differences due to race have not been studied. Because gabapentin is primarily renally excreted and there are no important racial differences in creatinine clearance, pharmacokinetic differences due to race are not expected.

Gabapentin pharmacokinetics were determined in 48 pediatric subjects between the ages of 1 month and 12 years following a dose of approximately were similar across the entire age group and occurred 2 to 3 hours postdose. In general, pediatric subjects between 1 month and <5 years of age achieved approximately 30% lower exposure (AUC) than that observed in those 5 years of age and older. Accordingly, oral clearance ormalized per body weight was higher in the younger children. Apparent oral clearance of gabapentin was directly proportional to creatinine clearance. Gabapentin elimination half-life averaged 4.7 hours and was similar across the age groups studied. A population pharmacokinetic analysis was performed in 253 pediatric subjects between 1 month

and 13 years of age. Patients received 10 to 65 mg/kg/day given three times a day. Apparen oral clearance (CL/F) was directly proportional to creatinine clearance and this relationship was similar following a single dose and at steady-state. Higher oral clearance values were observed in children <5 years of age compared to those observed in children 5 years of age and older, when normalized per body weight. The clearance was highly variable in infants <1 year of age. The normalized DLF values observed in pediatric patients 5 years of age and older were consistent with values observed in adults after a single dose. The oral volume of distribution normalized per body weight was constant across the age range.

These pharmacokinetic data indicate that the effective daily dose in pediatric patients with epilepsy ages 3 and 4 years should be 40 mg/kg/day to achieve average plasma concen similar to those achieved in patients 5 years of age and older receiving gabapentin at 30 mg/kg/day [see Dosage and Administration (2.2)].

Adult Patients with Renal Impairment

no effect on gabapentin pharmacokinetics.

Subjects (N=60) with renal impairment (mean creatinine clearance ranging from 13 to 114 mL min) were administered single 400 mg oral doses of gabapentin. The mean gabapentin half-life ranged from about 6.5 hours (patients with creatinine clearance >60 mL/min) to 52 hours (creatinine clearance <30 mL/min) and gabapentin renal clearance from about 90 mL/min (<60 mL/min group) to about 10 mL/min (<30 mL/min). Mean plasma clearance (CL/F) decreased from approximately 190 mL/min to 20 mL/min (see Dosage and Administration (2.3) and Use in Specific Populations (8.6)]. Pediatric patients with renal insufficiency have not been studied Hemodialysis

In a study in anuric adult subjects (N=11), the apparent elimination half-life of gabapentin on nondialysis days was about 132 hours; during dialysis the apparent half-life of gabapentin was reduced to 3.8 hours. Hemodialysis thus has a significant effect on gabapentin elimination in anuric subjects [see Dosage and Administration (2.3) and Use in Specific Populations (8.6)]. Hepatic Disease

Because gabapentin is not metabolized, no study was performed in patients with hepatic impairment.

Drug Interactions In Vitro Studies In vitro studies were conducted to investigate the potential of gabapentin to inhibit the major

cytochrome P450 enzymes (CYP1A2, CYP2A6, CYP2C9, CYP2C19, CYP2D6, CYP2E1, and CYP3A4) that mediate drug and xenobiotic metabolism using isoform selective marker substrates and human liver microsomal preparations. Only at the highest concentration tested (171 mcg mL; 1 mM) was a slight degree of inhibition (14% to 30%) of isoform CYP2A6 observed. No inhibition of any of the other isoforms tested was observed at gabapentin concentrations up to 171 mcg/mL (approximately 15 times the C_{max} at 3,600 mg/day). In Vivo Studies

The drug interaction data described in this section were obtained from studies involving healthy

adults and adult patients with epilepsy. In a single (400 mg) and multiple dose (400 mg three times a day) study of gabapentin in epileptic patients (N=8) maintained on phenytoin monotherapy for at least 2 months, gabapentin had no effect on the steady-state trough plasma concentrations of phenytoin and phenytoin had

Steady-state trough plasma carbamazepine and carbamazepine 10, 11 epoxide concentrations were not affected by concomitant gabapentin (400 mg three times a day; N=12) administration. Likewise, gabapentin pharmacokinetics were unaltered by carbamazepine adm

Valproic Acid The mean steady-state trough serum valproic acid concentrations prior to and during concomitan gabapentin administration (400 mg three times a day; N=17) were not different and neither were gabapentin pharmacokinetic parameters affected by valproic acid.

Phenobarbital Estimates of steady-state pharmacokinetic parameters for phenobarbital or gabapenting (300 mg three times a day; N=12) are identical whether the drugs are administered alone or

together.

Coadministration (N=18) of naproxen sodium capsules (250 mg) with gabapentin (125 mg) appears to increase the amount of gabapentin absorbed by 12% to 15%. Gabapentin had no effect on naproxen pharmacokinetic parameters. These doses are lower than the therapeutic doses for both drugs. The magnitude of interaction within the recommended dose ranges of either drug

Suite

5200

400 Connell Drive

Berkeley Heights,

NJ 07922

Coadministration of gabapentin (125 to 500 mg; N=48) decreases hydrocodone (10 mg; N=50) C_{max} and AUC values are 3% to 4% lower, respectively, after administration of hydrocodone alone; C_{max} and AUC values are 3% to 4% lower, respectively, after administration of 125 mg gabapentin and 21% to 22% lower, respectively, after administration of 500 mg gabapentin. The mechanism for this interaction is unknown. Hydrocodone increases gabapentin AUC values by 14%. The magnitude of interaction at other doses is not known

Morphine A literature article reported that when a 60 mg controlled-release morphine capsule was administered 2 hours prior to a 600 mg gabapentin capsule (N=12), mean gabapentin AUC increased by 44% compared to gabapentin administered without morphine. Morphine pharmacokinetic parameter values were not affected by administration of gabapentin 2 hours after morphine. The magnitude of interaction at other doses is not known

Cimetidine In the presence of cimetidine at 300 mg four times a day (N-12), the mean apparent oral clearance of gabapentin fell by 14% and creatinine clearance fell by 10%. Thus, cimetidine appeared to alter the renal excretion of both gabanentin and creatinine, an endogenous marker of appeared to ance the retail extension of our gasapentin and extension, an enoughous market of renal function. This small decrease in excretion of gasapentin by cimetidine is not expected to be of clinical importance. The effect of gasapentin on cimetidine was not evaluated.

Oral Contraceptive Based on AUC and half-life, multiple-dose pharmacokinetic profiles of norethindrone and ethinyl estradiol following administration of tablets containing 2.5 mg of norethindrone acetate and 50 meg of ethinyl estradiol were similar with and without coadministration of gabapentin (400 mg three times a day; N=13). The C_{max} of norethindrone was 13% higher when it was coadministered with gabapentin; this interaction is not expected to be of clinical importance

Antacid (Maalox®) containing magnesium and aluminum hydroxides reduced the mean bioavailability of gabapentin (N=16) by about 20%. This decrease in bioavailability was about 10% when gabapentin was administered 2 hours after Maalox. Probenecid Probenecid is a blocker of renal tubular secretion. Gabapentin pharmacokinetic parameters

without and with probenecid were comparable. This indicates that gabapentin does not undergo renal tubular secretion by the pathway that is blocked by probenecid. 13 NONCLINICAL TOXICOLOGY

Antacid (Maalox®) (aluminum hydroxide, magnesium hydroxide)

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility Carcinogenesis

Gabapentin was administered orally to mice and rats in 2-year carcinogenicity studies. No evidence of drug-related carcinogenicity was observed in mice treated at doses up to 2,000 mg/kg/day. At 2,000 mg/kg, the plasma gabapentin exposure (AUC) in mice was approximately 2 times that in humans at the MRHD of 3,600 mg/day. In rats, increases in the incidence of pancreatic acinar cell adenoma and carcinoma were found in male rats receiving the highest dose (2,000 mg/kg), but not at doses of 250 or 1,000 mg/kg/day. At 1,000 mg/kg, the plasma gabapentin exposure (AUC) in rats was approximately 5 times that in humans at the MRHD. Studies designed to investigate the mechanism of gabapentin-induced pancreatic carcinogenesis

in rats indicate that gabapentin stimulates DNA synthesis in rat pancreatic actinar cells in vitro and, thus, may be acting as a tumor promoter by enhancing mitogenic activity. It is not known whether gabapentin has the ability to increase cell proliferation in other cell types or in other

Gabapentin did not demonstrate mutagenic or genotoxic potential in *in vitro* (Ames test, HGPRT forward mutation assay in Chinese hamster lung cells) and *in vivo* (chromosomal aberration and micronucleus test in Chinese hamster hone marrow mouse micronucleus unscheduled DNA synthesis in rat hepatocytes) assays. Impairment of Fertility

No adverse effects on fertility or reproduction were observed in rats at doses up to 2,000 mg/kg. At 2,000 mg/kg, the plasma gabapentin exposure (AUC) in rats is approximately 8 times that in

14 CLINICAL STUDIES

14.1 Postherpetic Neuralgia Gabapentin was evaluated for the management of postherpetic neuralgia (PHN) in two randomized, double-blind, placebo-controlled, multicenter studies. The intent-to-treat (ITT) population consisted of a total of 563 patients with pain for more than 3 months after healing the herpes zoster skin rash (Table 6).

TABLE 6. Controlled PHN Studies: Duration, Dosages, and Number of Patien

Study	Study Duration	Gabapentin (mg/ day) ^a Target Dose	Patients Receiving Gabapentin	Patients Receiving Placebo
1	8 weeks	3,600	113	116
2	7 weeks	1,800, 2,400	223	111
		Total	336	227

a Given in 3 divided doses (TID)

Each study included a 7-or 8-week double-blind phase (3 or 4 weeks of titration and 4 weeks of fixed dose). Patients initiated treatment with titration to a maximum of 900 mg/day gabapentin over 3 days. Dosages were then to be titrated in 600 to 1,200 mg/day increments at 3- to 7-day intervals to the target dose over 3 to 4 weeks. Patients recorded their pain in a daily diary using an 11-point numeric pain rating scale ranging from 0 (no pain) to 10 (worst possible pain). A mean pain score during baseline of at least 4 was required for randomization. Analyses were conducted using the ITT population (all randomized patients who received at least one dose of study medication).

Both studies demonstrated efficacy compared to placebo at all doses tested. The reduction in weekly mean pain scores was seen by Week 1 in both studies, and were maintained to the end of treatment. Comparable treatment effects were observed in all active

Pharmacokinetic/pharmacodynamic modeling provided confirmatory evidence of efficacy across all doses. Figures 1 and 2 show pain intensity scores over time for Studies 1 and 2.Figure 1. Weekly Mean Pain Scores (Observed Cases in ITT Population): Study 1






Figure 2. Weekly Mean Pain Scores (Observed Cases in ITT Population): Study 2



The proportion of responders (those patients reporting at least 50% improve pain score compared to baseline) was calculated for each study (Figure 3). Figure 3. Proportion of Responders (patients with ≥50% reduction in pain score) at



The effectiveness of gabapentin as adjunctive therapy (added to other antiepileptic drugs) was tablished in multicenter placebo-controlled, double-blind, parallel-group clinical trials in adult and pediatric patients (3 years and older) with refractory partial seizures Evidence of effectiveness was obtained in three trials conducted in 705 patients (age 12 years

and above) and one trial conducted in 247 pediatric patients (3 to 12 years of age). The patients enrolled had a history of at least 4 partial seizures per month in spite of receiving one or more antiepileptic drugs at therapeutic levels and were observed on their established antiepileptic drug regimen during a 12-week baseline period (6 weeks in the study of pediatric patients). In patients continuing to have at least 2 (or 4 in some studies) seizures per month, gabapentin or placebo was then added on to the existing therapy during a 12-week treatment period. Effectiveness was assessed primarily on the basis of the percent of patients with a 50% or greater reduction in seizure frequency from baseline to treatment (the "responder rate") and a derived measure called response ratio, a measure of change defined as (T-B)/(T+B), in which B is the patient's baseline seizure frequency and T is the patient's seizure frequency during treatment. Response ratio is distributed within the range -1 to +1. A zero value indicates no change while complete elimination of seizures would give a value of -1; increased seizure rates would give a value of -1; increased seizure rates would give a value of seizure rates would give a value of seizure rates would give a value of -1; increased seizure rates would give positive values. A response ratio of -0.33 corresponds to a 50% reduction in seizure frequency. The results given below are for all partial seizures in the intent-to-treat (all patients who received any doses of treatment) population in each study, unless otherwise indicated.

One study compared gabapentin 1,200 mg/day, in three divided doses with placebo. Responder rate was 23% (14/61) in the gabapentin group and 9% (6/66) in the placebo group; the difference between groups was statistically significant. Response ratio was also better in the gabapentin group (-0.199) than in the placebo group (-0.044), a difference that also achieved statistical A second study compared primarily gabapentin 1,200 mg/day, in three divided doses (N=101),

with placebo (N=98). Additional smaller gabapentin dosage groups (600 mg/day, N=53; 1,800 mg/day, N=54) were also studied for information regarding dose response. Responder rate was higher in the gabapentin 1,200 mg/day group (16%) than in the placebo group (8%), but the difference was not statistically significant. The responder rate at 600 mg (17%) was also not significantly higher than in the placebo, but the responder rate in the 1,800 mg group (26%) was statistically significantly superior to the placebo rate. Response ratio was better in the gabapentin 1,200 mg/day group (-0.103) than in the placebo group (-0.022); but this difference was also not group (-0.105) and 1,800 mg/day group (-0.222) than in the 1,200 mg/day group, with the 1,800 mg/day group achieving statistical significance compared to the placebo group A third study compared gabapentin 900 mg/day, in three divided doses (N=111), and placebo

Medicines are

Call your doctor f FDA-1088.

medical

advice about

side

side

effects to FDA at 1-800

Store

provider or pharmacist.

all

the

possible

effects

gabapentin

tablets.

For

more

effect that bothers

 $^{\circ}$

that does not go away. information, ask your healthcare

provider if you

also statistically significantly superior in the gabapentin 900 mg/day group (-0.119) compared to that in the placebo group (-0.027), as was response ratio in 1,200 mg/day gabapentin (-0.184) compared to placebo. Analyses were also performed in each study to examine the effect of gabapentin on preventing secondarily generalized tonic-clonic seizures. Patients who experienced a secondarily generalized tonic-clonic seizures in either the baseline or in the treatment period in all three


(N=109). An additional gabapentin 1,200 mg/day dosage group (N=52) provided doseresponse data. A statistically significant difference in responder rate was seen in the gabapentin

900 mg/day group (22%) compared to that in the placebo group (10%). Response ratio was

placebo-controlled studies were included in these analyses. There were several response ratio comparisons that showed a statistically significant advantage for gabapentin compared to placebo and favorable trends for almost all comparisons.

Analysis of responder rate using combined data from all three studies and all doses (N=162, gabapentin, N=89, placebo) also showed a significant advantage for gabapentin over placebo in reducing the frequency of secondarily generalized tonic-clonic seizures.

In two of the three controlled studies, more than one dose of gabapentin was used. Within each study, the results did not show a consistently increased response to dose. However, looking across studies, a trend toward increasing efficacy with increasing dose is evident (see Figure 4). Figure 4. Responder Rate in Patients Receiving Gabapentin Expressed as a Difference from Placebo by Dose and Study: Adjunctive Therapy Studies in Patients ≥12 Years of Age with Partial Seizures



In the figure, treatment effect magnitude, measured on the Y axis in terms of the difference in the proportion of gabapentin and placebo-assigned patients attaining a 50% or greater reduction cure frequency from baseline, is plotted against the daily dose of gabapentin administered

Although no formal analysis by gender has been performed, estimates of response (Response Ratio) derived from clinical trials (398 men, 307 women) indicate no important gender differences exist. There was no consistent pattern indicating that age had any effect on the response to gabapentin. There were insufficient numbers of patients of races other than Caucasian to permit a comparison of efficacy among racial groups.

A fourth study in pediatric patients age 3 to 12 years compared 25 to 35 mg/kg/day gabapentin (N=118) with placebo (N=127). For all partial seizures in the intent-to-treat population, the response ratio was statistically significantly better for the gabapentin group (-0.146) than for the placebo group (-0.079). For the same population, the responder rate for gabapentin (21%) was not significantly different from placebo (18%). A study in pediatric patients age 1 month to 3 years compared 40 mg/kg/day gabapentin (N=38)

with placebo (N=38) in patients who were receiving at least one marketed antiepileptic drug and had at least one partial seizure during the screening period (within 2 weeks prior to baseline). Patients had up to 48 hours of baseline and up to 72 hours of double-blind video EEG monitoring to record and count the occurrence of seizures. There were no statistically significant differences between treatments in either the response ratio or responder rate.

16 HOW SUPPLIED/STORAGE AND HANDLING Gabapentin tablets. USP are supplied as follows:

Gabapentin tablets, USP 600 mg are White to off white, elliptical, biconvex film-coated tablets functionally scored on both sides, debossed with "C" and "5" on either side of score line on

one side.	
Bottles of 30's count:	NDC 42385-979-30
Bottles of 90's count:	NDC 42385-979-90
Bottles of 100's count:	NDC 42385-979-01
Bottles of 500's count:	NDC 42385-979-05
Bottles of 1,000's count:	NDC 42385-979-11
Carton with 100 (10 x 10) Unit-Dose Tablets	NDC 42385-979-72
containing PVC/PVDC blisters	

Gabapentin tablets, USP 800 mg are White to off white, elliptical, biconvex film-coated tablets functionally scored on both sides, debossed with "C" and "7" on either side of score line on

Bottles of 30's count NDC 42385-980-30 Bottles of 90's count: NDC 42385-980-90 NDC 42385-980-01 Bottles of 100's count: Bottles of 500's count: NDC 42385-980-05 NDC 42385-980-11 Bottles of 1,000's count Carton with 100 (10 x 10) Unit-Dose Tablets NDC 42385-980-72 containing PVC/PVDC blisters

Store at 20° to 25° C (68° to 77° F); excursions permitted between 15° to 30° C (59° to 86° F). [See USP Controlled Room Temperature.]

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide) Administration Information

Inform patients that gabapentin is taken orally with or without food. Inform patients that, should they divide the scored 600 mg or 800 mg tablet in order to administer a half-tablet, they should take the unused half-tablet as the next dose. Advise patients to discard half-tablets not used within 28 days of dividing the scored tablet.

Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Multiorgan Hypersensitivity Prior to initiation of treatment with gabapentin, instruct patients that a rash or other signs or symptoms of hypersensitivity (such as fever or lymphadenopathy) may herald a serious medical event and that the patient should report any such occurrence to a physician immediately [see Varnings and Precautions (5.1)]. Anaphylaxis and Angioedema

Advise patients to discontinue gabapentin and seek medical care if they develop signs or symptoms of anaphylaxis or angioedema [see Warnings and Precautions (5.2)]. Dizziness and Somnolence and Effects on Driving and Operating Heavy Machinery

Advise patients that gabapentin may cause dizziness, somnolence, and other symptoms and signs of CNS depression. Other drugs with sedative properties may increase these symptoms. Accordingly, although patients' ability to determine their level of impairment can be unreliable, advise them neither to drive a car nor to operate other complex machinery until they have gained sufficient experience on gabapentin to gauge whether or not it affects their mental and/or motor performance adversely. Inform patients that it is not known how long this effect lasts [see Warnings and Precautions (5.3) and Warnings and Precautions (5.4)]. Suicidal Thinking and Behavior

Counsel the patient, their caregivers, and families that AEDs, including gabapentin, may increase the risk of suicidal thoughts and behavior. Advise patients of the need to be alert for the emergence or worsening of symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts about self-harm. Instruct patients to report behaviors of concern immediately to healthcare providers [see Warnings and Precautions (5.6)].

Respiratory Depression

Inform patients about the risk of respiratory depression. Include information that the risk is greatest for those using concomitant CNS depressants (such as opioid analgesics) or those with underlying respiratory impairment. Teach patients how to recognize respiratory depression and advise them to seek medical attention immediately if it occurs [see Warnings and Precautions (5.7)1.Use in Pregnancy

Instruct patients to notify their physician if they become pregnant or intend to become pregnant

during therapy, and to notify their physician if they are breast feeding or intend to breast feed during therapy [see Use in Specific Populations (8.1) and (8.2)]. The brands listed are trademarks of their respective owners and are not trademarks of Laurus

Dispense with Medication Guide available at: $\underline{https://www.laurusgenerics.us/gab\text{-}tabs\text{-}medi\text{-}guide.pdf}$

Manufactured for: Laurus Generics Inc 400 Connell Drive Suite 5200 Berkeley Heights, NJ 07922 Manufactured by:

Laurus Labs Limited Anakapalli-531011

(gab" a pen' tin) What is the most important information I should know about gabapentin tablets?

Do not stop taking gabapentin tablets without first talking to your healthcare provider. Stopping gabapentin tablets suddenly can cause serious problems Gabapentin tablets can cause serious side effects including:

MEDICATION GUIDE

Gabapentin Tablets, USF

1. Suicidal Thoughts. Like other antiepileptic drugs, gabapentin tablets may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.

Call a healthcare provider right away if you have any of these symptoms, especially if the

thoughts about suicide or dying attempts to commit suicide

new or worse depression

new or worse anxiety feeling agitated or restless

panic attacks trouble sleeping (insomnia) new or worse irritability

difficulty with speaking

usually of legs and

unusual double

eye movement

difficulty with jerky movements

coordination

vision

lack of coordination most common

side effects

gabapentin

tablets include:

drowsy and vomiting

, which could

increase the occurrence

of

accidental

injury, including falls

taking gabapentin tablets?

I should know about gabapentin tab blets. See "What I should avoid while

lets?"

gabapentin

serious side effects including:

infection

Call your healthcare provider between visits as needed, especially if you are worried about

Do not stop taking gabapentin tablets without first talking to a healthcare provider. • Stopping gabapentin tablets suddenly can cause serious problems. Stopping a seizure medicine suddenly in a patient who has epilepsy can cause seizures that will not stop (status

epilepticus). Suicidal thoughts or actions can be caused by things other than medicines. If you have

suicidal thoughts or actions, your healthcare provider may check for other causes.

2. Changes in behavior and thinking - Using gabapentin tablets in children 3 to 12 years of age can cause emotional changes, aggressive behavior, problems with concentration, rechanges in school performance, and hyperactivity.

3. Gabapentin tablets may cause serious or life-threatening allergic reactions that may affect your skin or other parts of your body such as your liver or blood cells. This may cause you to be hospitalized or to stop gabapentin tablets. You may or may not have a rash with an allergic reaction caused by gabapentin tablets. Call a healthcare provider right away if you have any of

the following symptoms skin rash

· difficulty breathing

 fever swollen glands that do not go away

swelling of your face, lips, throat, or tongue

yellowing of your skin or of the whites of the eyes

unusual bruising or bleeding severe fatigue or weakness unexpected muscle pain

age and older with seizures.

• frequent infections

4. Serious breathing problems. Serious breathing problems can occur when gabapentin tablets are taken with other medicines that can cause severe sleepiness or decreased awareness, or when it is taken by someone who already has breathing problems. Watch for increased sleep

These symptoms may be the first signs of a serious reaction. A healthcare provider should examine you to decide if you should continue taking gabapentin tablets.

decreased breathing when starting gabapentin tablets or when the dose is increased. Get help right away if breathing problems occur. What are gabapentin tablets? Gabapentin tablets are a prescription medicine used to treat:

 Pain from damaged nerves (postherpetic pain) that follows healing of shingles (a painful rash that comes after a herpes zoster infection) in adults. Partial seizures when taken together with other medicines in adults and children 3 years of

Who should not take gabapentin tablets? Do not take gabapentin tablets if you are allergic to gabapentin or any of the other ingredients in gabapentin tablets. See the end of this Medication Guide for a complete list of ingredients in

gabapentin tablets. What should I tell my healthcare provider before taking gabapentin tablets?

Before taking gabapentin tablets, tell your healthcare provider if you: have or have had kidney problems or are on hemodialysis

have or have had depression, mood problems, or suicidal thoughts or behavior

have diabetes

have breathing problems are pregnant or plan to become pregnant. It is not known if gabapentin tablets can harm your unborn baby. Tell your healthcare provider right away if you become pregnant while taking gabapentin tablets. You and your healthcare provider will decide if you should take

are breast-feeding or plan to breast-feed. Gabapentin can pass into breast milk. You and your healthcare provider should decide how you will feed your baby while you take gabapentin Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take any opioid pain medicine (such as oxycodone), any medicines for anxiety

(such as lorazepam) or insomnia (such as zolpidem), or any medicines that make you sleepy. You may have a higher chance for dizziness, sleepiness, or breathing problems if these medicines are taken with gabapentin tablets. Taking gabapentin tablets with certain other medicines can cause side effects or affect how well they work. Do not start or stop other medicines without talking to your healthcare provider

pharmacist when you get a new medicine How should I take gabapentin tablets?

What should I avoid while taking gabapentin tablets?

What are the possible side effects of gabapentin tablets?

gabapentin tablets?

· lack of coordination

viral infection

feeling tired

gabapentin tablets while you are pregnant.

• Take gabapentin tablets exactly as prescribed. Your healthcare provider will tell you how many gabapentin tablets to take.

Know the medicines you take. Keep a list of them and show it to your healthcare provider and

• Do not change your dose of gabapentin tablets without talking to your healthcare provider. If you take gabapentin tablets and break a tablet in half, the unused half of the tablet should be taken at your next scheduled dose. Half tablets not used within 28 days of breaking should be thrown away.

Gabapentin tablets can be taken with or without food. If you take an antacid containing aluminum and magnesium, such as Maalox®, Mylanta®, Gelusil®, Gaviscon®, or Di-Gel®, you should wait at least 2 hours before taking your next dose of gabapentin tablets. you take too many gabapentin tablets, call your healthcare provider or your local Poison Control Center right away at 1-800-222-1222.

Do not drink alcohol or take other medicines that make you sleepy or dizzy while taking gabapentin tablets without first talking with your healthcare provider. Taking gabapentin tablets with alcohol or drugs that cause sleepiness or dizziness may make your or dizziness worse. • Do not drive, operate heavy machinery, or do other dangerous activities until you know how gabapentin tablets affects you. Gabapentin tablets can slow your thinking and motor skills

Gabapentin tablets may cause serious side effects including: See "What is the most important information I should know about gabapentin tablets?" problems driving while using gabapentin tablets. See "What I should avoid while taking

sleepiness and dizziness, which could increase the occurrence of accidental injury, including • The most common side effects of gabapentin tablets include:

· feeling drowsy

nausea and vomiting

· difficulty with speaking jerky movements difficulty with coordination tremor · swelling, usually of legs and feet double vision

 fever Tell your healthcare provider if you have any side effect that bothers you or that does not go

These are not all the possible side effects of gabapentin tablets. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA

at 1-800-FDA-1088. How should I store gabapentin tablets? Store gabapentin tablets between 68°F to 77°F (20°C to 25°C).

Keep gabapentin tablets and all medicines out of the reach of children.

General information about the safe and effective use of gabapentin tablets Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use gabapentin tablets for a condition for which it was not prescribed. Do not give gabapentin tablets to other people, even if they have the same symptoms that you have. It may

you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about gabapentin tablets that was written for healthcare

For more information, call Laurus Generics Inc. at 1-833-3-LAURUS (1-833-352-8787).

What are the ingredients in gabapentin tablets? Active ingredient: gabapentin USP Inactive ingredients: copovidone, maize starch 5%, microcrystalline cellulose, magnesium stearate and poloxamer P407. In addition, the film coating contains the following inactive

ingredients: Polyethylene Glycol 3350, Polyvinyl alcohol-part hydrolyzed, titanium oxide and This Medication Guide has been approved by the U.S. Food and Drug Administration

The brands listed are trademarks of their respective owners and are not trademarks of Laurus Dispense with Medication Guide available at

 $\underline{https://www.laurusgenerics.us/gab\text{-}tabs\text{-}medi\text{-}guide.pdf}$ Manufactured for: Laurus Generics Inc. 400 Connell Drive Suite 5200

Berkeley Heights, NJ 07922 Manufactured by: Laurus Labs Limited

Anakapalli-531011 Revised: 10/2023

Do not Taking

with gabapentin tablets

Know

the medicines you take.

when you

get a new medicine

tablets to

gabapentin tablets

exactly

as

}€------

Manufactured by:
Laurus Labs Limited India Revised: 10/2023 Anakapalli-531011

Manufactured https://www.laurusgenerics.us/gab-tabs-medi-guide.pdf Laurus Generics l

for:

poloxamer Pa Glycol 3350, Dispense with Medication Guide available at: The brands listed This Medication Guide has Polyvinyl are trademarks

of their respective owners

trademarks of Laurus Labs Limited

Inactive ingredients: poloxamer P407. In a

Active ingredient: gabapentin What are the ingredients in like more information, talk with your healthcare provider. For more information, call Laurus Generics Inc. about Guide summarizes addition, gabapentin same gabapentin USP tablets the at 5%, Ħ

(1-833-352

-8787).

profession

lthc

care provider or pharmacist als.

gabapentin tablets. If you would

dition, the film coating alcohol-part hydrolyzed, by the U.S. g contains the fo l, titanium oxide a Food nicrocrystalline of the following inactive ingredients: Polyethylene magnesium

stearate and

gabapentin tablets for a conceptople, even if they have the General information about the safe and sometimes prescribed for purposes for which it was no symptoms that you most important information about was other than those 1-833-3-LAURUS not prescribed. You can ask your heal

This Medication

for information

Keep gabapentin tablets and all medicines out of the reach How should I store gabapentin tablets? gabapentin tablets between 68°F to 77°F effective use have. It may harm them. of gabapentin of children. listed Do tablets ed in not a Me t give Me gabapentin tablets to other dication Guide. Do not use

acting aggressive, being angry, or violent acting on dangerous impuls · an extreme increase in activity and talking (mania) · other unusual changes in behavior or mood How can I watch for early symptoms of suicidal thoughts and actions? · Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or · Keep all follow-up visits with your healthcare provider as scheduled

problems driving while using sleepiness and dizziness, whi

See Gabapentin tablets may cause What are the possible side effects of gabapentin tablets? tablets "What is the most important information

20

may make your

operate heavy machinery,

 10

do other

dangerous

s worse.

vities until

notor skills.

you know how gabapentin

 10

. Gabapentin tablets

can slow your

thinking

and

motor

right away you

What should I avoid while taking 1 take too first talking

before taking your next dose e too many gabapentin at 1-800-222-1222. tablets, , call

of

Do not drink alcohol or take other medicines that make you sleepy with your healthcare provider. Taking tablets? gabapentin y or dizzy while pentin tablets

your healthcare provider or your le taking gabapentin tablets with alcohol or drugs that

local Poison Control Center

gabapentin

gabapentin

Gaviscon' an au -Gel® , you containing aluminum and should wait at least 2 hours

antacid

Do not change your dose of gabapentin tablets without gabapentin tablets taken Half tablets not used and with break $^{\circ}$ a tablet within food. in half, the \mathbf{H} 28 days of

Gabapentin tablets

be

such as Maalox®

your next scheduled dose.

If you take

talking to provider will tell care provider.

unused sed half of t f breaking s half should be thrown away. the tablet should be taken a

How should I take gabapentin tablets? entin tablets with certain other medicines stop other medicines without talking to prescribed. Your healthcare sleepiness, sleepy. to y and your healthcare or breathing to problems effects or Ħ affect how well they work re provider and pharmacist you how many gabapentin these medicines are taken

as zolpidem), or

You may have a higher vitamins, and herbal supplements. (such provider as oxycodone), all the medicines any medicines Especially

for dizziness, tell your healthc including aslor prescription and over-the-